

**NEET SPECIAL**

**NEET Phase-1 Solved Paper 2016**

(More than 60% Questions from MTG Books / Magazines)

**NEET Phase-2 Practice Paper**

# CHEMISTRY

# today

**India's #1**  
CHEMISTRY MONTHLY FOR  
JEE (Main & Advanced) & NEET

Solved Papers 2016

- **Karnataka CET**
- **Kerala PET**

**NCERT CORNER**

**ACE**  
YOUR WAY **CBSE XII**  
SERIES 1

**EXAM PREP**

**CONCEPT  
MAP**

**mtg**

Trust of more than  
1 Crore Readers  
Since 1982



**CHEMISTRY  
MUSING**

**OLYMPIAD  
PROBLEMS**

**You Ask ?**

**We Answer ✓**

# CHEMISTRY today

Volume 25

No. 6

June 2016

**Managing Editor**  
Mahabir Singh

**Editor**  
Anil Ahlawat  
(BE, MBA)

**Corporate Office:**

Plot 99, Sector 44 Institutional area, Gurgaon - 122 003 (HR).

Tel : 0124-6601200 e-mail : info@mtg.in website : www.mtg.in

**Regd. Office:**

406, Taj Apartment, Near Safdarjung Hospital, New Delhi - 110029.

## CONTENTS

Chemistry Musing Problem Set 35	8
NEET Phase I Solved Paper 2016	10
NEET Phase II Practice Paper 2016	18
Concept Booster	26
NCERT Corner	31
Exam Prep	41
Concept Map	46
Olympiad Problems	50
Ace Your Way CBSE XII	54
Kerala PET Solved Paper 2016	63
Karnataka CET Solved Paper 2016	70
Advanced Chemistry Bloc	77
You Ask, We Answer	80
Chemistry Musing Solution Set 34	83
Crossword	85



### TO OUR READERS

We are happy that intelligent students, teachers and other professionals continue to patronise Mathematics Today, Chemistry Today, Physics For You and Biology Today.

To them, we are addressing this open letter in view of increase in the cost of production and postage in the last five years. All round spiralling prices have pushed production costs so high, that many in our fraternity find it impossible to continue business. We are compelled to raise the price to ₹ 40 from July 2016 issue.

We understand the pressure of cost on the student-teacher community in general but, we are hoping our readers will understand our problems and that we have no option but to comply with this unavoidable move.

We on our part, will keep up our efforts to improve the magazines in all its aspects.

Subscribe online at [www.mtg.in](http://www.mtg.in)

Individual Subscription Rates				Combined Subscription Rates			
	1 yr.	2 yrs.	3 yrs.		1 yr.	2 yrs.	3 yrs.
Mathematics Today	330	600	775	PCM	900	1500	1900
Chemistry Today	330	600	775	PCB	900	1500	1900
Physics For You	330	600	775	PCMB	1000	1800	2300
Biology Today	330	600	775				

Send D.D/M.O in favour of MTG Learning Media (P) Ltd.

Payments should be made directly to : MTG Learning Media (P) Ltd,

Plot No. 99, Sector 44, Gurgaon - 122003 (Haryana)

We have not appointed any subscription agent.

Owned, Printed and Published by Mahabir Singh from 406, Taj Apartment, New Delhi - 29 and printed by Personal Graphics and Advertisers (P) Ltd., Okhla Industrial Area, Phase-II, New Delhi. Readers are advised to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/Infocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only.

Editor : Anil Ahlawat

Copyright© MTG Learning Media (P) Ltd.

All rights reserved. Reproduction in any form is prohibited.

# CHEMISTRY MUSING

**C**hemistry Musing was started from August '13 issue of Chemistry Today with the suggestion of Shri Mahabir Singh. The aim of Chemistry Musing is to augment the chances of bright students preparing for JEE (Main and Advanced) / NEET / AIIMS / PETs with additional study material.

In every issue of Chemistry Today, 10 challenging problems are proposed in various topics of JEE (Main and Advanced) / NEET. The detailed solutions of these problems will be published in next issue of Chemistry Today.


The readers who have solved five or more problems may send their solutions. The names of those who send atleast five correct solutions will be published in the next issue.

We hope that our readers will enrich their problem solving skills through "Chemistry Musing" and stand in better stead while facing the competitive exams.



## PROBLEM Set 35

### JEE MAIN/NEET

1. A balloon filled with helium rises to a certain height at which it gets fully inflated to a volume of  $1 \times 10^5$  litres. If at this altitude temperature and atmospheric pressure are 268 K and  $2 \times 10^{-3}$  atm respectively, what weight of helium will be required to fully inflate the balloon?  
(a) 24.62 g (b) 0.35 g  
(c) 8.97 g (d) 36.36 g
2. The enthalpy of neutralisation of a weak acid in 1 M solution with a strong base is  $-56.1 \text{ kJ mol}^{-1}$ . If enthalpy of ionisation of the acid is  $1.5 \text{ kJ mol}^{-1}$  and enthalpy of neutralisation of the strong acid with a strong base is  $-57.3 \text{ kJ mol}^{-1}$ , what is the % ionisation of the weak acid in molar solution (Assume the acid to be monobasic)?  
(a) 10 (b) 15 (c) 20 (d) 25
3. Salicylic acid is treated with bromine under two different conditions as shown :



Compounds X and Y in the given reactions are respectively



4. Which of the following does not react with AgCl?  
(a)  $\text{Na}_2\text{S}_2\text{O}_3$  (b)  $\text{NH}_4\text{OH}$   
(c)  $\text{NaNO}_3$  (d)  $\text{Na}_2\text{CO}_3$
5. Identify the number of optically active isomers of  $\text{C}_3\text{H}_4\text{ClBr}$ .  
(a) 2 (b) 4 (c) 3 (d) 5


### JEE ADVANCED

6. A 5.00 g sample of a natural gas, consisting of methane ( $\text{CH}_4$ ) and ethylene ( $\text{C}_2\text{H}_4$ ) was burnt in excess oxygen, yielding 14.5 g of  $\text{CO}_2$  and some  $\text{H}_2\text{O}$  as products. What is the percentage of ethylene in the sample?  
(a) 20.2% (b) 40.4% (c) 10.1% (d) 38.8%



## COMPREHENSION

When a quaternary ammonium halide is treated with moist silver oxide (a slurry of  $\text{Ag}_2\text{O}$  in water), a quaternary ammonium hydroxide is formed as the main product. Thermal decomposition of the quaternary ammonium hydroxide gives an alkene. This reaction is known as Hofmann elimination and has most of the characteristics of an  $\text{E}_2$  reaction.



7. The Hofmann elimination reactions are  
 (i) concerted  
 (ii) stereoselective for *anti* elimination  
 (iii) the reactions that occur in separate steps  
 (iv) stereoselective for *syn* elimination.  
 (a) (i) and (iv) (b) (i) and (ii)  
 (c) only (i) (d) only (iii).
8. In the given reaction, which of the following is correct regarding yield of X and Y?

- (a)  $X = Y$  (b)  $X \approx Y$   
 (c)  $X > Y$  (d)  $X < Y$

## INTEGER VALUE

9. The number of moles of sodium acetate that should be dissolved to prepare one litre of 0.063 molar buffer solution of pH 4.5 is  $x \times 10^{-2}$ . The value of  $x$  is
10. Two 1.0 g carbon disks 1.00 cm apart have opposite charges of equal magnitude such that there is  $1.00 \times 10^{-5}$  N force between them. The ratio of excess electrons to total atoms on the negatively charged disk is nearly  $x \times 10^{-14}$  electron/atom. The value of  $x$  is

## Solution Senders of Chemistry Musing

### SET-33

- |                     |                |
|---------------------|----------------|
| 1. G. Geeth Nischal | Andhra Pradesh |
| 2. Samrat Gupta     | Kolkata        |

### SET-34

- |                 |             |
|-----------------|-------------|
| 1. Indranil Roy | West Bengal |
|-----------------|-------------|

## ATTENTION COACHING INSTITUTES: a great offer from MTG

**MTG** offers "Classroom Study Material" for JEE (Main & Advanced), NEET and FOUNDATION MATERIAL for Class 7, 8, 9, 10, 11 & 12 with YOUR BRAND NAME & COVER DESIGN.

This study material will save you lots of money spent on teachers, typing, proof-reading and printing. Also, you will save enormous time. Normally, a good study material takes 2 years to develop. But you can have the material printed with your logo delivered at your doorstep.

Profit from associating with MTG Brand – the most popular name in educational publishing for JEE (Main & Advanced)/NEET/PMT ....

Order sample chapters on Phone/Fax/e-mail.


Phone : 0124-6601200

09312680856

e-mail : sales@mtg.in | www.mtg.in



## CLASSROOM STUDY MATERIAL



Your logo  
here



# NEET

Phase-I  
held on 1<sup>st</sup>  
May

## SOLVED PAPER 2016

Hurray!!

We are happy to inform our readers that out of the 45 questions asked in NEET 2016, more than 60% questions were either exactly same or of similar type from the **MTG Books**.

Hurray!!

Here, the references of few are given :

S. No.	MTG Books	Q. No.	Pg. No.
2	NCERT Fingertips	93	151
4	NEET Guide	138	550
5	NCERT Fingertips	97	125
7	NEET Guide	127	159
8	NEET Guide	72	621
20	NEET Guide	194	517
21	NCERT Fingertips	35	313
22	Objective Chemistry	164	766

S. No.	MTG Books	Q. No.	Pg. No.
23	Objective Chemistry	91	31
26	NEET Guide	90	594
30	NEET Guide	43	751
32	NCERT Fingertips	36	64
40	Objective Chemistry	82	203
41	NCERT Fingertips	80	149
43	NEET Guide	74	849
44	NEET Guide	35	8

and more such questions .....

- Which of the following statements about the composition of the vapour over an ideal 1 : 1 molar mixture of benzene and toluene is correct? Assume that the temperature is constant at 25°C. (Given, vapour pressure data at 25°C, benzene = 12.8 kPa, toluene = 3.85 kPa)
  - The vapour will contain equal amounts of benzene and toluene.
  - Not enough information is given to make a prediction.
  - The vapour will contain a higher percentage of benzene.
  - The vapour will contain a higher percentage of toluene.
- Match the compounds given in column I with the hybridisation and shape given in column II and mark the correct option.

	Column I		Column II
(A)	XeF <sub>6</sub>	(i)	Distorted octahedral
(B)	XeO <sub>3</sub>	(ii)	Square planar
(C)	XeOF <sub>4</sub>	(iii)	Pyramidal
(D)	XeF <sub>4</sub>	(iv)	Square pyramidal

Code :

- |     |          |          |          |          |
|-----|----------|----------|----------|----------|
|     | <b>A</b> | <b>B</b> | <b>C</b> | <b>D</b> |
| (a) | (iv)     | (iii)    | (i)      | (ii)     |
| (b) | (iv)     | (i)      | (ii)     | (iii)    |
| (c) | (i)      | (iii)    | (iv)     | (ii)     |
| (d) | (i)      | (ii)     | (iv)     | (iii)    |
- The correct statement regarding the comparison of staggered and eclipsed conformations of ethane, is
    - the eclipsed conformation of ethane is more stable than staggered conformation even though the eclipsed conformation has torsional strain
    - the staggered conformation of ethane is more stable than eclipsed conformation, because staggered conformation has no torsional strain
    - the staggered conformation of ethane is less stable than eclipsed conformation, because staggered conformation has torsional strain
    - the eclipsed conformation of ethane is more stable than staggered conformation, because eclipsed conformation has no torsional strain.

4. Fog is a colloidal solution of  
 (a) solid in gas (b) gas in gas  
 (c) liquid in gas (d) gas in liquid.

5. Match items of Column I with the items of Column II and assign the correct code :

	Column I		Column II
(A)	Cyanide process	(i)	Ultrapure Ge
(B)	Froth floatation process	(ii)	Dressing of ZnS
(C)	Electrolytic reduction	(iii)	Extraction of Al
(D)	Zone refining	(iv)	Extraction of Au
		(v)	Purification of Ni

Code :

**A    B    C    D**

- (a) (i)    (ii)    (iii)    (iv)  
 (b) (iii)    (iv)    (v)    (i)  
 (c) (iv)    (ii)    (iii)    (i)  
 (d) (ii)    (iii)    (i)    (v)
6. Which is the correct statement for the given acids?  
 (a) Phosphinic acid is a monoprotic acid while phosphonic acid is a diprotic acid.  
 (b) Phosphinic acid is a diprotic acid while phosphonic acid is a monoprotic acid.  
 (c) Both are diprotic acids.  
 (d) Both are triprotic acids.
7. The correct thermodynamic conditions for the spontaneous reaction at all temperatures is  
 (a)  $\Delta H < 0$  and  $\Delta S > 0$  (b)  $\Delta H < 0$  and  $\Delta S < 0$   
 (c)  $\Delta H < 0$  and  $\Delta S = 0$  (d)  $\Delta H > 0$  and  $\Delta S < 0$
8. Which one of the following statements is correct when  $\text{SO}_2$  is passed through acidified  $\text{K}_2\text{Cr}_2\text{O}_7$  solution?  
 (a)  $\text{SO}_2$  is reduced.  
 (b) Green  $\text{Cr}_2(\text{SO}_4)_3$  is formed.  
 (c) The solution turns blue.  
 (d) The solution is decolourised.
9. The correct statement regarding RNA and DNA, respectively is  
 (a) the sugar component in RNA is a arabinose and the sugar component in DNA is ribose  
 (b) the sugar component in RNA is 2'-deoxyribose and the sugar component in DNA is arabinose  
 (c) the sugar component in RNA is arabinose and the sugar component in DNA is 2'-deoxyribose  
 (d) the sugar component in RNA is ribose and the sugar component in DNA is 2'-deoxyribose.

10. Which of the following reagents would distinguish *cis*-cyclopenta-1-2-diol from the *trans*-isomer?

(a)  $\text{MnO}_2$  (b) Aluminium isopropoxide  
 (c) Acetone (d) Ozone

11. The correct statement regarding a carbonyl compound with a hydrogen atom on its alpha-carbon, is

(a) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as carbonylation  
 (b) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as keto-enol tautomerism  
 (c) a carbonyl compound with a hydrogen atom on its alpha-carbon never equilibrates with its corresponding enol  
 (d) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as aldehyde-ketone equilibration.

12. At  $100^\circ\text{C}$  the vapour pressure of a solution of 6.5 g of a solute in 100 g water is 732 mm. If  $K_b = 0.52$ , the boiling point of this solution will be

(a)  $102^\circ\text{C}$  (b)  $103^\circ\text{C}$   
 (c)  $101^\circ\text{C}$  (d)  $100^\circ\text{C}$

13. Consider the nitration of benzene using mixed conc.  $\text{H}_2\text{SO}_4$  and  $\text{HNO}_3$ . If a large amount of  $\text{KHSO}_4$  is added to the mixture, the rate of nitration will be

(a) unchanged (b) doubled  
 (c) faster (d) slower.

14. The pressure of  $\text{H}_2$  required to make the potential of  $\text{H}_2$ -electrode zero in pure water at 298 K is

(a)  $10^{-10}$  atm (b)  $10^{-4}$  atm  
 (c)  $10^{-14}$  atm (d)  $10^{-12}$  atm

15. The correct statement regarding the basicity of arylamines is

(a) arylamines are generally more basic than alkylamines because of aryl group  
 (b) arylamines are generally more basic than alkylamines, because the nitrogen atom in arylamines is *sp*-hybridised  
 (c) arylamines are generally less basic than alkylamines because the nitrogen lone-pair electrons are delocalised by interaction with the aromatic ring  $\pi$ -electron system

- (d) arylamines are generally more basic than alkylamines because the nitrogen lone-pair electrons are not delocalised by interaction with the aromatic ring  $\pi$ -electron system.
16. In a protein molecule various amino acids are linked together by  
 (a) peptide bond (b) dative bond  
 (c)  $\alpha$ -glycosidic bond (d)  $\beta$ -glycosidic bond.
17. Consider the molecules  $\text{CH}_4$ ,  $\text{NH}_3$  and  $\text{H}_2\text{O}$ . Which of the given statements is false?  
 (a) The  $\text{H}-\text{O}-\text{H}$  bond angle in  $\text{H}_2\text{O}$  is smaller than the  $\text{H}-\text{N}-\text{H}$  bond angle in  $\text{NH}_3$ .  
 (b) The  $\text{H}-\text{C}-\text{H}$  bond angle in  $\text{CH}_4$  is larger than the  $\text{H}-\text{N}-\text{H}$  bond angle in  $\text{NH}_3$ .  
 (c) The  $\text{H}-\text{C}-\text{H}$  bond angle in  $\text{CH}_4$ , the  $\text{H}-\text{N}-\text{H}$  bond angle in  $\text{NH}_3$ , and the  $\text{H}-\text{O}-\text{H}$  bond angle in  $\text{H}_2\text{O}$  are all greater than  $90^\circ$ .  
 (d) The  $\text{H}-\text{O}-\text{H}$  bond angle in  $\text{H}_2\text{O}$  is larger than the  $\text{H}-\text{C}-\text{H}$  bond angle in  $\text{CH}_4$ .
18. Which of the following statements is false?  
 (a)  $\text{Ca}^{2+}$  ions are not important in maintaining the regular beating of the heart.  
 (b)  $\text{Mg}^{2+}$  ions are important in the green parts of the plants.  
 (c)  $\text{Mg}^{2+}$  ions form a complex with ATP.  
 (d)  $\text{Ca}^{2+}$  ions are important in blood clotting.
19. Which one of the following orders is correct for the bond dissociation enthalpy of halogen molecules?  
 (a)  $\text{Br}_2 > \text{I}_2 > \text{F}_2 > \text{Cl}_2$  (b)  $\text{F}_2 > \text{Cl}_2 > \text{Br}_2 > \text{I}_2$   
 (c)  $\text{I}_2 > \text{Br}_2 > \text{Cl}_2 > \text{F}_2$  (d)  $\text{Cl}_2 > \text{Br}_2 > \text{F}_2 > \text{I}_2$
20. The rate of first-order reaction is  $0.04 \text{ mol L}^{-1} \text{ s}^{-1}$  at 10 seconds and  $0.03 \text{ mol L}^{-1} \text{ s}^{-1}$  at 20 seconds after initiation of the reaction. The half-life period of the reaction is  
 (a) 44.1 s (b) 54.1 s  
 (c) 24.1 s (d) 34.1 s
21. Which one given below is a non-reducing sugar?  
 (a) Glucose (b) Sucrose  
 (c) Maltose (d) Lactose
22. Which one of the following characteristics is associated with adsorption?  
 (a)  $\Delta G$  and  $\Delta H$  are negative but  $\Delta S$  is positive.  
 (b)  $\Delta G$  and  $\Delta S$  are negative but  $\Delta H$  is positive.  
 (c)  $\Delta G$  is negative but  $\Delta H$  and  $\Delta S$  are positive.  
 (d)  $\Delta G$ ,  $\Delta H$  and  $\Delta S$  all are negative.
23. Two electrons occupying the same orbital are distinguished by  
 (a) azimuthal quantum number  
 (b) spin quantum number  
 (c) principal quantum number  
 (d) magnetic quantum number.
24. Lithium has a *bcc* structure. Its density is  $530 \text{ kg m}^{-3}$  and its atomic mass is  $6.94 \text{ g mol}^{-1}$ . Calculate the edge length of a unit cell of lithium metal. ( $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$ )  
 (a) 527 pm (b) 264 pm  
 (c) 154 pm (d) 352 pm
25. The pair of electrons in the given carbanion,  $\text{CH}_3\text{C}\equiv\text{C}^-$ , is present in which of the following orbitals?  
 (a)  $sp^2$  (b)  $sp$   
 (c)  $2p$  (d)  $sp^3$
26. The product obtained as a result of a reaction of nitrogen with  $\text{CaC}_2$  is  
 (a)  $\text{CaCN}_3$  (b)  $\text{Ca}_2\text{CN}$   
 (c)  $\text{Ca}(\text{CN})_2$  (d)  $\text{CaCN}$
27. In the reaction  

$$\text{H}-\text{C}\equiv\text{CH} \xrightarrow[\text{(ii) CH}_3\text{CH}_2\text{Br}]{\text{(i) NaNH}_2/\text{liq. NH}_3} \text{X} \xrightarrow[\text{(ii) CH}_3\text{CH}_2\text{Br}]{\text{(i) NaNH}_2/\text{liq. NH}_3} \text{Y}$$
  
 X and Y are  
 (a) X = 2-butyne, Y = 2-hexyne  
 (b) X = 1-butyne, Y = 2-hexyne  
 (c) X = 1-butyne, Y = 3-hexyne  
 (d) X = 2-butyne, Y = 3-hexyne.
28. *MY* and *NY*<sub>3</sub>, two nearly insoluble salts, have the same  $K_{sp}$  values of  $6.2 \times 10^{-13}$  at room temperature. Which statement would be true in regard to *MY* and *NY*<sub>3</sub>?  
 (a) The salts *MY* and *NY*<sub>3</sub> are more soluble in 0.5 M *KY* than in pure water.  
 (b) The addition of the salt of *KY* to solution of *MY* and *NY*<sub>3</sub> will have no effect on their solubilities.  
 (c) The molar solubilities of *MY* and *NY*<sub>3</sub> in water are identical.  
 (d) The molar solubility of *MY* in water is less than that of *NY*<sub>3</sub>.
29. When copper is heated with conc.  $\text{HNO}_3$  it produces  
 (a)  $\text{Cu}(\text{NO}_3)_2$ ,  $\text{NO}$  and  $\text{NO}_2$   
 (b)  $\text{Cu}(\text{NO}_3)_2$  and  $\text{N}_2\text{O}$   
 (c)  $\text{Cu}(\text{NO}_3)_2$  and  $\text{NO}_2$   
 (d)  $\text{Cu}(\text{NO}_3)_2$  and  $\text{NO}$




30. The product formed by the reaction of an aldehyde with a primary amine is  
 (a) carboxylic acid (b) aromatic acid  
 (c) Schiff's base (d) ketone.

31. The addition of a catalyst during a chemical reaction alters which of the following quantities?  
 (a) Enthalpy (b) Activation energy  
 (c) Entropy (d) Internal energy

32. Predict the correct order among the following :  
 (a) bond pair - bond pair > lone pair - bond pair > lone pair - lone pair  
 (b) lone pair - bond pair > bond pair - bond pair > lone pair - lone pair  
 (c) lone pair - lone pair > lone pair - bond pair > bond pair - bond pair  
 (d) lone pair - lone pair > bond pair - bond pair > lone pair - bond pair


33. Consider the following liquid-vapour equilibrium.



Which of the following relations is correct?

- (a)  $\frac{d \ln P}{dT^2} = \frac{-\Delta H_v}{T^2}$  (b)  $\frac{d \ln P}{dT} = \frac{\Delta H_v}{RT^2}$   
 (c)  $\frac{d \ln G}{dT^2} = \frac{\Delta H_v}{RT^2}$  (d)  $\frac{d \ln P}{dT} = \frac{-\Delta H_v}{RT}$

34. Which of the following biphenyls is optically active?




35. Which of the following statements about hydrogen is incorrect?

- (a) Hydronium ion,  $\text{H}_3\text{O}^+$  exists freely in solution.  
 (b) Dihydrogen does not act as a reducing agent.  
 (c) Hydrogen has three isotopes of which tritium is the most common.  
 (d) Hydrogen never acts as cation in ionic salts.

36. The electronic configurations of Eu (Atomic No. 63), Gd (Atomic No. 64) and Tb (Atomic No. 65) are

- (a)  $[\text{Xe}]4f^6 5d^1 6s^2$ ,  $[\text{Xe}]4f^7 5d^1 6s^2$  and  $[\text{Xe}]4f^8 5d^1 6s^2$   
 (b)  $[\text{Xe}]4f^7 6s^2$ ,  $[\text{Xe}]4f^7 5d^1 6s^2$  and  $[\text{Xe}]4f^9 6s^2$   
 (c)  $[\text{Xe}]4f^7 6s^2$ ,  $[\text{Xe}]4f^8 6s^2$  and  $[\text{Xe}]4f^8 5d^1 6s^2$   
 (d)  $[\text{Xe}]4f^6 5d^1 6s^2$ ,  $[\text{Xe}]4f^7 5d^1 6s^2$  and  $[\text{Xe}]4f^9 6s^2$


37. The reaction



can be classified as

- (a) dehydration reaction  
 (b) Williamson alcohol synthesis reaction  
 (c) Williamson ether synthesis reaction  
 (d) alcohol formation reaction.

38. For the following reactions :



Which of the following statements is correct?

- (a) (A) is elimination, (B) and (C) are substitution reactions.  
 (b) (A) is substitution, (B) and (C) are addition reactions.  
 (c) (A) and (B) are elimination reactions and (C) is addition reaction.  
 (d) (A) is elimination, (B) is substitution and (C) is addition reaction.

39. In which of the following options the order of arrangement does not agree with the variation of property indicated against it?
- I < Br < Cl < F (increasing electron gain enthalpy)
  - Li < Na < K < Rb (increasing metallic radius)
  - Al<sup>3+</sup> < Mg<sup>2+</sup> < Na<sup>+</sup> < F<sup>-</sup> (increasing ionic size)
  - B < C < N < O (increasing first ionisation enthalpy)
40. Equal moles of hydrogen and oxygen gases are placed in a container with a pin-hole through which both can escape. What fraction of the oxygen escapes in the time required for one-half of the hydrogen to escape?
- 3/8
  - 1/2
  - 1/8
  - 1/4
41. Among the following, the correct order of acidity is
- HClO<sub>2</sub> < HClO < HClO<sub>3</sub> < HClO<sub>4</sub>
  - HClO<sub>4</sub> < HClO<sub>2</sub> < HClO < HClO<sub>3</sub>
  - HClO<sub>3</sub> < HClO<sub>4</sub> < HClO<sub>2</sub> < HClO
  - HClO < HClO<sub>2</sub> < HClO<sub>3</sub> < HClO<sub>4</sub>
42. Which of the following is an analgesic?
- Streptomycin
  - Chloromycetin
  - Novalgin
  - Penicillin
43. Natural rubber has
- alternate *cis*- and *trans*-configuration
  - random *cis*- and *trans*-configuration
  - all *cis*-configuration
  - all *trans*-configuration.
44. The ionic radii of A<sup>+</sup> and B<sup>-</sup> ions are 0.98 × 10<sup>-10</sup> m and 1.81 × 10<sup>-10</sup> m. The coordination number of each ion in AB is
- 8
  - 2
  - 6
  - 4
45. Which of the following has longest C—O bond length? (Free C—O bond length in CO is 1.128 Å.)
- [Fe(CO)<sub>4</sub>]<sup>2-</sup>
  - [Mn(CO)<sub>6</sub>]<sup>+</sup>
  - Ni(CO)<sub>4</sub>
  - [Co(CO)<sub>4</sub>]<sup>-</sup>


### SOLUTIONS

1. (c) :  $p_{\text{Benzene}} = x_{\text{Benzene}} p_{\text{Benzene}}^{\circ}$   
 $p_{\text{Toluene}} = x_{\text{Toluene}} p_{\text{Toluene}}^{\circ}$   
 For an ideal 1 : 1 molar mixture of benzene and toluene,  $x_{\text{Benzene}} = \frac{1}{2}$  and  $x_{\text{Toluene}} = \frac{1}{2}$   
 $p_{\text{Benzene}} = \frac{1}{2} p_{\text{Benzene}}^{\circ} = \frac{1}{2} \times 12.8 \text{ kPa} = 6.4 \text{ kPa}$


$$p_{\text{Toluene}} = \frac{1}{2} p_{\text{Toluene}}^{\circ} = \frac{1}{2} \times 3.85 \text{ kPa} = 1.925 \text{ kPa}$$

Thus, the vapour will contain a high percentage of benzene as the partial vapour pressure of benzene is higher as compared to that of toluene.

2. (c) :



3. (b) :




Newman's projections of ethane

Magnitude of torsional strain depends upon the angle of rotation about C—C bond. Staggered form has the least torsional strain and the eclipsed form has the maximum torsional strain so, the staggered conformation of ethane is more stable than the eclipsed conformation.

4. (c) : Fog is an example of aerosol in which dispersed phase is liquid and dispersion medium is gas.
5. (c) : Mac-Arthur Forest cyanide process is used for the extraction of gold and silver.
- Froth floatation process is used for the concentration of sulphide ores *e.g.*, ZnS.
- Electrolytic reduction is used for the extraction of highly electropositive metals like Na, K, Mg, Ca, Al, etc.
- Zone refining method is used for producing semiconductor and other metals of very high purity *e.g.*, Ge, Si, B, Ga and In. Purification of Ni is done by vapour phase refining (Mond's process).

6. (a):



7. (a, c):  $\Delta G = \Delta H - T\Delta S$

If  $\Delta H < 0$  and  $\Delta S > 0$

$\Delta G = (-ve) - T(+ve)$

then at all temperatures,  $\Delta G = -ve$ , spontaneous reaction.


If  $\Delta H < 0$  and  $\Delta S = 0$

$\Delta G = (-ve) - T(0) = -ve$  at all temperatures.

8. (b):  $K_2Cr_2O_7 + H_2SO_4 + 3SO_2 \rightarrow K_2SO_4 + Cr_2(SO_4)_3 + H_2O$   
(Green)


9. (d)

10. (c):



*Trans*-isomer does not react with acetone.

11. (b): Keto-enol tautomerism:



12. (c): Given:  $W_B = 6.5$  g,  $W_A = 100$  g,  $p_s = 732$  mm,  $K_b = 0.52$ ,  $T_b^\circ = 100^\circ\text{C}$ ,  $p^\circ = 760$  mm

$$\frac{p^\circ - p_s}{p^\circ} = \frac{n_2}{n_1} \Rightarrow \frac{760 - 732}{760} = \frac{n_2}{100/18}$$

$$\Rightarrow n_2 = \frac{28 \times 100}{760 \times 18} = 0.2046 \text{ moles}$$


$$\Delta T_b = K_b \times m$$

$$T_b - T_b^\circ = K_b \times \frac{n_2 \times 1000}{W_A(\text{g})}$$

$$T_b - 100^\circ\text{C} = \frac{0.52 \times 0.2046 \times 1000}{100} = 1.06$$

$$T_b = 100 + 1.06 = 101.06^\circ\text{C}$$


13. (d): Mechanism of nitration is:



If a large amount of  $KHSO_4$  is added then conc. of  $HSO_4^-$  ions increases and the reaction will be shifted in backward direction hence, the rate of nitration will be slower.

14. (c): pH = 7 for water.

$$-\log[H^+] = 7 \Rightarrow [H^+] = 10^{-7}$$




$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.0591}{2} \log \frac{p_{H_2}}{[H^+]^2}$$

$$0 = 0 - \frac{0.0591}{2} \log \frac{p_{H_2}}{(10^{-7})^2}$$

$$\log \frac{p_{H_2}}{(10^{-7})^2} = 0 \Rightarrow \frac{p_{H_2}}{(10^{-7})^2} = 1 \quad [\because \log 1 = 0]$$

$$p_{H_2} = 10^{-14} \text{ atm}$$


15. (c):



In arylamines, lone pair of electrons on nitrogen atom is delocalised over the benzene ring, thus, not available for donation. So, arylamines are less basic than alkylamines ( $R-\ddot{N}H_2$ ).

16. (a)

17. (d):



18. (a):  $Ca^{2+}$  ions are required to trigger the contraction of muscles and to maintain the regular beating of the heart.


### Your favourite MTG Books/Magazines available in WEST BENGAL at

- Progressive Book Centre-Kharagpur, Ph: 03222-279956
- Rukmani Agencies-Kolkata, Ph: 033-24666173, 0224839473
- Katha-O-Kahani Book Seller Pvt. Ltd.-Kolkata, Ph: 033-22196313
- International Book Trust-Kolkata, Ph: 033-22414947, 24792343
- Novelty Books-Siliguri, Ph: 0353-2525445
- Om Traders-Siliguri, Ph: 0353-6450299, 09749048104

Visit "MTG IN YOUR CITY" on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call 0124-6601200 for further assistance.









Hydrogen has three isotopes of which protium is the most common and tritium is radioactive.

36. (b)


37. (c): Williamson's ether synthesis reaction involves the treatment of sodium alkoxide with a suitable alkyl halide to form an ether.



Saturated compound is converted into unsaturated compound by removal of group of atoms hence, it is an elimination reaction.



—Br group is replaced by —OH group hence, it is a substitution reaction.



Addition of  $\text{Br}_2$  converts an unsaturated compound into a saturated compound hence, it is an addition reaction.

39. (a, d): The correct order of increasing negative electron gain enthalpy is:  $\text{I} < \text{Br} < \text{F} < \text{Cl}$  and the correct order of increasing first ionisation enthalpy is  $\text{B} < \text{C} < \text{O} < \text{N}$ .


40. (c): Let the number of moles of each gas =  $x$

Fraction of hydrogen escaped =  $\frac{1}{2}x$

$$\begin{aligned} \frac{r_{\text{O}_2}}{r_{\text{H}_2}} &= \sqrt{\frac{M_{\text{H}_2}}{M_{\text{O}_2}}} \\ \Rightarrow \frac{n_{\text{O}_2}/t}{\frac{x}{2}/t} &= \sqrt{\frac{2}{32}} = \sqrt{\frac{1}{16}} = \frac{1}{4} \\ \Rightarrow \frac{n_{\text{O}_2}/t}{\frac{x}{2}/t} &= \frac{1}{4} \Rightarrow n_{\text{O}_2} = \frac{1}{8}x \end{aligned}$$

Hence, fraction of oxygen escaped =  $\frac{1}{8}$

41. (d): The acidic character of the oxoacids increases with increase in oxidation number of the halogen atom *i.e.*,



This can be explained on the basis of relative stability of the anions left after removal of a proton. Since

the stability of the anion decreases in the order:

$\text{ClO}_4^- > \text{ClO}_3^- > \text{ClO}_2^- > \text{ClO}^-$ , acid strength also decreases in the same order.

42. (c): Streptomycin, chloromycetin and penicillin are antibiotics while novalgin is an analgesic.

43. (c): Natural rubber is *cis*-polyisoprene.

44. (c): Radius ratio,  $\frac{r_+}{r_-} = \frac{0.98 \times 10^{-10}}{1.81 \times 10^{-10}} = 0.541$

It lies in the range of 0.414 to 0.732 hence, coordination number of each ion will be 6 as the compound will have NaCl type structure *i.e.*, octahedral arrangement.

45. (a): The greater the negative charge on the carbonyl complex, the more easy it would be for the metal to permit its electrons to participate in the back bonding, the higher would be the  $M-C$  bond order and simultaneously there would be larger reduction in the  $C-O$  bond order. Thus,  $[\text{Fe}(\text{CO})_4]^{2-}$  has the lowest  $C-O$  bond order means the longest bond length.



### Your favourite MTG Books/Magazines available in ASSAM & NORTH EAST at

- **Books Corner- Agartala**  
Ph: 0381-2301945; Mob: 985635894
- **Suravi Sahitya Kutir- Dibrugarh**  
Ph: 0373-2322085; Mob: 9435002641, 9435031727
- **Prime Books & Periodicals- Guwahati**  
Ph: 0361-2733065; Mob: 9954126701
- **Book Emporium- Guwahati**  
Ph: 0361-2635094; Mob: 9864057226
- **Books View- Guwahati**  
Mob: 9706081383
- **Jain Book Shop- Imphal**  
Mob: 9856031157
- **PC Jain- Imphal**  
Ph: 0388-2451756, 2225004; Mob: 9856084649
- **Naveen Pustakalaya- Jorhat**  
Ph: 0376-2323940; Mob: 9435514204, 9435051093
- **Abhan- Silchar**  
Ph: 03842-231844
- **Haque Book Stall- Tezpur**  
Ph: 03712-231129; Mob: 9435081257

Visit "MTG IN YOUR CITY" on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call **0124-6601200** for further assistance.



# BOOST your **NEET** score

## Practice paper for phase II

- On combustion in excess of air, lithium forms mainly the oxide, sodium forms peroxide while potassium or rubidium or caesium forms superoxides. It suggests that as cationic size increases the
  - anionic polarisability increases, thus stability decreases
  - anionic size increases, polarisability increases, covalency increases, thus stability increases
  - anionic size increases, coordination number increases, lattice energy increases, thus stability increases
  - anionic size decreases, stability increases.
- The reaction of  $\text{KMnO}_4$  and  $\text{HCl}$  results in
  - oxidation of Mn in  $\text{KMnO}_4$  and production of  $\text{Cl}_2$
  - reduction of Mn in  $\text{KMnO}_4$  and production of  $\text{H}_2$
  - oxidation of Mn in  $\text{KMnO}_4$  and production of  $\text{H}_2$
  - reduction of Mn in  $\text{KMnO}_4$  and production of  $\text{Cl}_2$ .
- The ether that undergoes electrophilic substitution reaction is
  - $\text{CH}_3\text{OC}_2\text{H}_5$
  - $\text{C}_6\text{H}_5\text{OCH}_3$
  - $\text{CH}_3\text{OCH}_3$
  - $\text{C}_2\text{H}_5\text{OC}_2\text{H}_5$
- The boiling point of methanol is greater than methyl thiol because
  - there is intramolecular hydrogen bonding in methanol and intermolecular hydrogen bonding in methyl thiol
  - there is intermolecular hydrogen bonding in methanol and no hydrogen bonding in methyl thiol
  - there is no hydrogen bonding in methanol and intermolecular hydrogen bonding in methyl thiol
  - there is no hydrogen bonding in methanol and intramolecular hydrogen bonding in methyl thiol.
- For a reaction  $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$ , rate and rate constant are  $1.02 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$  and  $3.4 \times 10^{-5} \text{ s}^{-1}$ . The concentration of  $\text{N}_2\text{O}_5$  at that time will be
  - $1.732 \text{ mol L}^{-1}$
  - $3 \text{ mol L}^{-1}$
  - $1.02 \times 10^{-4} \text{ mol L}^{-1}$
  - $3.2 \times 10^5 \text{ mol L}^{-1}$
- With respect to benzyl carbocation and benzyl carbanion, the correct statement is
  - $\text{C}_6\text{H}_5\text{CH}_2^+$  ion is more stable than  $\text{CH}_3^+$  ion and  $\text{C}_6\text{H}_5\text{CH}_2^-$  ion is also more stable than  $\text{CH}_3^-$  ion
  - $\text{C}_6\text{H}_5\text{CH}_2^+$  ion is less stable than  $\text{CH}_3^+$  ion and  $\text{C}_6\text{H}_5\text{CH}_2^-$  ion is more stable than  $\text{CH}_3^-$  ion
  - $\text{C}_6\text{H}_5\text{CH}_2^+$  ion is more stable than  $\text{CH}_3^+$  ion and  $\text{C}_6\text{H}_5\text{CH}_2^-$  ion is less stable than  $\text{CH}_3^-$  ion
  - $\text{C}_6\text{H}_5\text{CH}_2^+$  ion is less stable than  $\text{CH}_3^+$  ion and  $\text{C}_6\text{H}_5\text{CH}_2^-$  ion is also less stable than  $\text{CH}_3^-$  ion.
- A cylinder contains 100 g of an ideal gas (mol. wt.  $40 \text{ g mol}^{-1}$ ) at  $27^\circ\text{C}$  and 2 atm pressure. In transference process, the cylinder fell and a dent was created, i.e., there was a decrease in the volume of cylinder. But valve attached to the cylinder cannot keep the pressure greater than 2 atm, so 10 g of gas leaked out. Calculate the volume of the cylinder after dent.
  - 3.08 L
  - 30.8 L
  - 2.770 L
  - 27.71 L
- A clathrate may be defined as a
  - cage compound
  - liquid crystal
  - mixture
  - solid solution.
- For the reaction,
$$\text{H}_{2(g)} + \text{CO}_{2(g)} \rightleftharpoons \text{CO}_{(g)} + \text{H}_2\text{O}_{(g)}$$



if the initial pressure of  $H_2 = CO_2$  and  $x$  moles/litre of hydrogen is consumed at equilibrium, the correct expression of  $K_p$  is

- (a)  $\frac{x^2}{1-x^2}$  (b)  $\frac{(1+x)^2}{(1-x)^2}$   
 (c)  $\frac{x^2}{(1-x)^2}$  (d)  $\frac{x^2}{(2+x)^2}$




10. Which of the following sets contains only addition polymers?

- (a) Polyethylene, polypropylene, terylene  
 (b) Polyethylene, PVC, acrilan  
 (c) Buna-S, nylon, polybutadiene  
 (d) Bakelite, PVC, polyethylene

11. The wave number of first line of Balmer series of hydrogen is  $15200\text{ cm}^{-1}$ . The wave number of the first Balmer line of  $Li^{2+}$  ion is

- (a)  $15200\text{ cm}^{-1}$  (b)  $60800\text{ cm}^{-1}$   
 (c)  $76000\text{ cm}^{-1}$  (d)  $136800\text{ cm}^{-1}$

12. Compound X on oxidation with alkaline  $KMnO_4$  gives benzoic acid. X may be

- (a)  (b)   
 (c)  (d) all of these.

13. Match list I with list II and select the correct answer using the codes given below the lists.

List I	List II
(Compound)	(Oxidation state of nitrogen)
(P) $N_2O_5$	(i) -2
(Q) $NaN_3$	(ii) +5
(R) NO	(iii) -1/3
(S) $N_2H_4$	(iv) +2

- P Q R S  
 (a) (ii) (iii) (iv) (i)  
 (b) (i) (ii) (iii) (iv)  
 (c) (iv) (i) (ii) (iii)  
 (d) (iii) (i) (iv) (ii)

14. Aldehydes and ketones cannot be distinguished by

- (a) Molisch's test (b) Tollens' test  
 (c) Benedict's test (d) Schiff's test.

15. The correct order of increasing ionic character of the given compounds is

- (a)  $BeCl_2 < MgCl_2 < CaCl_2 < BaCl_2$   
 (b)  $BeCl_2 < MgCl_2 < BaCl_2 < CaCl_2$   
 (c)  $BeCl_2 < BaCl_2 < MgCl_2 < CaCl_2$   
 (d)  $BaCl_2 < CaCl_2 < MgCl_2 < BeCl_2$

16. Which of the following is incorrect?

- (a) Molecular weight of NaCl found by osmotic pressure measurements is half of the theoretical value.  
 (b) Molecular weight of  $CH_3COOH$  in benzene found by cryoscopic methods is double of the theoretical value.  
 (c) Osmotic pressure of 0.1 M glucose solution is half of that of 0.1 M NaCl solution.  
 (d) Molecular weight of HCl found by any colligative property will be same in the aqueous solution and benzene solution.

17. Match list I (reagents used with ethyne) with list II (products) and select the correct answer using the codes given below the lists.

List I	List II
P. Hydrogen in presence of catalyst	(i) Benzene
Q. Heat at $600^\circ\text{C}$	(ii) Acetaldehyde
R. Hydrogen in presence of Pd and $CaCO_3$ at 473 K	(iii) Ethene
S. Water in presence of $H_2SO_4$ and $HgSO_4$	(iv) Ethane

- P Q R S  
 (a) (ii) (iii) (iv) (i)  
 (b) (i) (iv) (iii) (ii)  
 (c) (i) (iv) (ii) (iii)  
 (d) (iv) (i) (iii) (ii)

18. The sequence in which different reagents are used to separate each component in a mixture of PbS, ZnS and FeS from froth floatation process is

- (a) potassium ethyl xanthate, KCN  
 (b) potassium ethyl xanthate, KCN, NaOH,  $CuSO_4$ , acid  
 (c) KCN,  $CuSO_4$ , acid  
 (d) none of these.

19. Correct statement about carbonyl group is

- (a) it is non-planar  
 (b) carbon atom is  $sp^2$  hybridised  
 (c) oxygen has five non-bonding electrons  
 (d) carbon oxygen bond is non-polar.


20. Which of the following will produce hard water?

- (a) Saturation of water with  $\text{CaSO}_4$
- (b) Addition of  $\text{Na}_2\text{SO}_4$  to water
- (c) Saturation of water with  $\text{CaCO}_3$
- (d) Saturation of water with  $\text{MgCO}_3$

21. Which of the following sets consists only essential amino acids?

- (a) Leucine, Proline, glycine
- (b) Leucine, Histidine, Lysine
- (c) Valine, Glycine, Proline
- (d) Tyrosine, Arginine, Serine.

22. Consider the graph between compressibility factor  $Z$  and pressure  $P$



The correct increasing order of ease of liquefaction of the gases shown in the above graph is

- (a)  $\text{H}_2 < \text{N}_2 < \text{CH}_4 < \text{CO}_2$
- (b)  $\text{CO}_2 < \text{CH}_4 < \text{N}_2 < \text{H}_2$
- (c)  $\text{H}_2 < \text{CH}_4 < \text{N}_2 < \text{CO}_2$
- (d)  $\text{CH}_4 < \text{H}_2 < \text{N}_2 < \text{CO}_2$

23. Under identical conditions of temperature and pressure, the ratio of the rates of effusion of  $\text{O}_2$  and  $\text{CO}_2$  gases is given by

- (a)  $\frac{\text{rate of effusion of oxygen}}{\text{rate of effusion of carbon dioxide}} = 0.87$
- (b)  $\frac{\text{rate of effusion of oxygen}}{\text{rate of effusion of carbon dioxide}} = 1.17$
- (c)  $\frac{\text{rate of effusion of oxygen}}{\text{rate of effusion of carbon dioxide}} = 8.7$
- (d)  $\frac{\text{rate of effusion of oxygen}}{\text{rate of effusion of carbon dioxide}} = 11.7$


24. Which of the following statements is false?

- (a) Lower the concentration of D.O., more polluted is the water sample.
- (b) The tolerable limit of lead in drinking water is 50 ppb.
- (c) Water is considered pure if it has BOD less than 5 ppm.
- (d) In COD determination, the pollutants resistant to microbial oxidation are not oxidised by oxidising agent like  $\text{K}_2\text{Cr}_2\text{O}_7$ .

25. The number of electrons involved in the reaction when one Faraday of electricity is passed through an electrolyte in solution is

- (a)  $12 \times 10^{46}$
- (b) 96540
- (c)  $8 \times 10^{16}$
- (d)  $6 \times 10^{23}$


26. The reaction,



can be described as

- (a) acid-base reaction followed by an intramolecular  $\text{S}_{\text{N}}1$  reaction
- (b) acid-base reaction followed by an intramolecular Williamson's ether synthesis
- (c) E2 reaction followed by an addition reaction to a double bond
- (d)  $\text{S}_{\text{N}}2$  reaction following an intramolecular Williamson's ether synthesis.

27. The two complexes given below are



- (a) geometrical isomers
- (b) position isomers
- (c) optical isomers
- (d) identical.

28. Tertiary nitro compounds do not tautomerise because

- (a) there is no double bond
- (b) there is no  $\alpha$ -hydrogen
- (c) oxygen is more electronegative than hydrogen
- (d) all of the above.

29. The prefixes *syn* and *anti* are used to denote

- (a) structural isomers
- (b) conformational isomers
- (c) geometrical isomers
- (d) optical isomers.

30. Which of the following has correct increasing basic strength?

- (a)  $\text{MgO} < \text{BeO} < \text{CaO} < \text{BaO}$
- (b)  $\text{BeO} < \text{MgO} < \text{CaO} < \text{BaO}$
- (c)  $\text{BaO} < \text{CaO} < \text{MgO} < \text{BeO}$
- (d)  $\text{CaO} < \text{BaO} < \text{BeO} < \text{MgO}$


31. In a ferromagnetic material

- (a) all the magnetic moment vectors are aligned in one direction
- (b) half of the magnetic moment vectors point in one direction and rest in the opposite direction
- (c) all the magnetic moment vectors are randomly oriented
- (d) it is characterised by small magnetic moment.

32. The heats of neutralization of  $\text{CH}_3\text{COOH}$ ,  $\text{HCOOH}$ ,  $\text{HCN}$  and  $\text{H}_2\text{S}$  are  $-13.2$ ,  $-13.4$ ,  $-2.9$  and  $-3.8$  kcal per equivalent respectively. The correct increasing order of acid strength is

(a)  $\text{HCOOH} < \text{CH}_3\text{COOH} < \text{H}_2\text{S} < \text{HCN}$   
 (b)  $\text{HCN} < \text{H}_2\text{S} < \text{CH}_3\text{COOH} < \text{HCOOH}$   
 (c)  $\text{HCOOH} < \text{CH}_3\text{COOH} < \text{HCN} < \text{H}_2\text{S}$   
 (d)  $\text{CH}_3\text{COOH} < \text{H}_2\text{S} < \text{HCN} < \text{HCOOH}$

33. If the bond enthalpy of  $\text{HCl}_{(g)}$ ,  $\text{H}_{2(g)}$  and  $\text{Cl}_{2(g)}$  is 103, 104 and 58 kcal  $\text{mol}^{-1}$ , then the enthalpy change at constant volume for the following reaction at 300 K will be






(a)  $-44$  kcal  $\text{mol}^{-1}$  (b)  $-66$  kcal  $\text{mol}^{-1}$   
 (c)  $-11$  kcal  $\text{mol}^{-1}$  (d)  $-22$  kcal  $\text{mol}^{-1}$



34. In context with the transition elements, which of the following statements is incorrect?

(a) In addition to the normal oxidation states, the zero oxidation state is also shown by these elements in complexes.  
 (b) In the highest oxidation states, transition metals show basic character and form cationic complexes.  
 (c) In the highest oxidation states of the first five transition elements (Sc to Mn), all the 4s and 3d electrons are used for bonding.  
 (d) Once the  $d^5$  configuration is exceeded, the tendency to involve all the 3d electrons in bonding decreases.

35. Which of the following observations are correct about the major products X and Y of the following reaction?



(i) X is  (ii) X is 

(iii) Y is  (iv) Y is 

(a) (i) and (ii) (b) (ii) and (iv)  
 (c) (i) and (iii) (d) (i) and (iv)

36. Which of the following is most basic?

(a)  $\text{Ce}(\text{OH})_3$  (b)  $\text{Lu}(\text{OH})_3$   
 (c)  $\text{Yb}(\text{OH})_3$  (d)  $\text{Tb}(\text{OH})_3$

37. The correct statement about orthoboric acid is

(a) it is a strong monobasic acid  
 (b) it is not a proton donor, but a weak Lewis acid  
 (c) it is a tribasic acid  
 (d) it is harmful for eyes.

38. When phenol is treated with  $\text{Br}_2$  water, the product is

(a) *o*- and *p*-bromophenol  
 (b) 2,3,4-tribromophenol  
 (c) 2,4,6-tribromophenol  
 (d) none of these.


39. Arsenic drugs are mainly used in the treatment of

(a) jaundice (b) typhoid  
 (c) syphilis (d) cholera.

40. In the standardization of  $\text{Na}_2\text{S}_2\text{O}_3$  using  $\text{K}_2\text{Cr}_2\text{O}_7$  by iodometry, the equivalent weight of  $\text{K}_2\text{Cr}_2\text{O}_7$  is

(a) (molecular weight)/2  
 (b) (molecular weight)/6  
 (c) (molecular weight)/3  
 (d) same as the molecular weight.

41. Which of the following represents physical adsorption?



42. Secondary amine with Hinsberg's reagent forms

(a) *N*-alkyl sulphonamide soluble in KOH solution.  
 (b) *N*-alkyl sulphonamide insoluble in KOH solution.  
 (c) *N,N*-dialkyl sulphonamide soluble in KOH solution.  
 (d) *N,N*-dialkyl sulphonamide insoluble in KOH solution.



43. Primary alcohols can be prepared by the reaction of RMgX with  
 (a)  $C_2H_5CHO$  (b)  $CH_3CHO$   
 (c)  $HCHO$  (d)  $C_3H_7CHO$
44. Which one of the following statements is not true?  
 (a) The conjugate base of  $H_2PO_4^-$  is  $HPO_4^{2-}$ .  
 (b)  $pH + pOH = 14$  for all aqueous solutions.  
 (c) The pH of  $1 \times 10^{-8}$  M HCl is 8.  
 (d) pH decreases with rise in temperature.
45. Which is not the correct statement for ionic solids in which positive and negative ions are held by strong electrostatic attractive forces?  
 (a) The radius ratio  $r_+/r_-$  increases as coordination number increases.  
 (b) As the difference in size of ions increases, coordination number increases.  
 (c) When coordination number is eight,  $r_+/r_-$  ratio lies between 0.225 to 0.414.  
 (d) In ZnS type structure, anions have *ccp* arrangement and cations occupy alternate tetrahedral voids. the coordination number of  $Zn^{2+}$  and  $S^{2-}$  respectively are 4 and 4.

### SOLUTIONS

1. (c): The stability of the peroxides and superoxides increases as the metal ions become larger. Basically, larger cations are stabilised by larger anions because if both ions are similar in size, the coordination number will be high and this gives a high lattice energy.
2. (d):  $2KMnO_4 + 16HCl \longrightarrow 2KCl + 2MnCl_2 + 8H_2O + 5Cl_2$   
 In the reaction, manganese is reduced and changes from  $Mn^{+7}$  to  $Mn^{2+}$  while  $Cl^-$  is oxidised to  $Cl_2$ .
3. (b): Alkyl aryl ethers undergo electrophilic substitution reactions.
4. (b): Intermolecular H-bonding is present in methanol while there is no H-bonding in methyl thiol,  $CH_3SH$ .  

$$\begin{array}{ccccccc} \cdots & H-O & \cdots & H-O & \cdots & H-O & \cdots & H-O & \cdots \\ & | & & | & & | & & | & \\ & CH_3 & & CH_3 & & CH_3 & & CH_3 & \end{array}$$
 Methanol
5. (b): Unit of rate constant shows that it is a first order reaction.  
 $\therefore \text{Rate} = \text{Rate constant} [N_2O_5]$   

$$\Rightarrow [N_2O_5] = \frac{\text{Rate}}{\text{Rate constant}} = \frac{1.02 \times 10^{-4}}{3.4 \times 10^{-5}} = 3 \text{ M}$$
6. (a): Both +ve and -ve charge are delocalised on the benzene ring hence,  $C_6H_5^+\bar{C}H_2$  and  $C_6H_5^-\bar{C}H_2$ , both are stable.
7. (d): As 10 g of gas leaked out after the dent, 90 g of the gas must occupy the volume of the cylinder at 2 atm and  $27^\circ C$ .  

$$PV = \frac{w}{M} RT,$$

$$2 \times V = \frac{90}{40} \times 0.0821 \times 300 \Rightarrow V = 27.7 \text{ L}$$
8. (a): Clathrates are defined as cage compounds in which gaseous atoms are entrapped in cavities of crystal lattices of other compounds.
9. (c):  $H_{2(g)} + CO_{2(g)} \rightleftharpoons CO_{(g)} + H_2O_{(g)}$   
 Initial pressure      1                      1                      0                      0  
 At equilibrium (1 - x)                      (1 - x)                      x                      x  

$$K_p = \frac{P_{CO} \times P_{H_2O}}{P_{H_2} \times P_{CO_2}} = \frac{(x)(x)}{(1-x)(1-x)} = \frac{x^2}{(1-x)^2}$$
10. (b)
11. (d): Applying  $\bar{v} = R \left[ \frac{1}{n_1^2} - \frac{1}{n_2^2} \right] Z^2$   

$$\bar{v}_{Li^{2+}} = \bar{v}_H \times 3^2 = 15200 \times 9 = 136800 \text{ cm}^{-1}$$
12. (d): The alkyl side chain on benzene, irrespective of its size or nature, is completely oxidised to carboxylic group on oxidation with alkaline  $KMnO_4$ .
13. (a)
14. (a): Aldehydes and ketones cannot be distinguished by Molisch's test.
15. (a): Ionic character of compounds of metals increases down the group.
16. (d): In aqueous solution, HCl dissociates but in benzene it does not.
17. (d)
18. (b): Potassium ethyl xanthate acts as a collector, KCN and NaOH depress the froth floatation property of ZnS and FeS particles. Thus, only PbS go into froth. Then copper sulphate is added to the tank (mixture) which activates floating character of ZnS and only ZnS comes along with froth. The remaining slurry is acidified and FeS floats along with froth.
19. (b)
20. (a):  $CaSO_4$  is soluble in water and provides  $Ca^{2+}$  ions to develop hardness.  $CaCO_3$  and  $MgCO_3$  are insoluble in water.


21. (b)

22. (a):  $Z$  is less than 1 means more compressible gases while  $Z > 1$  means less compressible gases.

23. (b):  $\frac{r_{O_2}}{r_{CO_2}} = \sqrt{\frac{M_{CO_2}}{M_{O_2}}} = \sqrt{\frac{44}{32}} = 1.17$

24. (d): In COD determination, the pollutants resistant to microbial oxidation are also oxidised by oxidising agent like  $K_2Cr_2O_7$ .


25. (d): One Faraday of electricity means one mole of electrons i.e.,  $6 \times 10^{23}$  electrons.




27. (d): Both represent only one molecule and no isomerism is seen.

28. (b): Tertiary nitro compounds do not have any  $\alpha$ -hydrogen atom hence, do not show tautomerism.

29. (c): *Syn* and *anti* prefixes are used to denote geometrical isomers involving either  $C=N$  or  $N=N$  bond, e.g.,




30. (b): The basic strength of the oxides increases gradually as the ionization enthalpy of metal decreases down the group.



31. (a): In ferromagnetic materials, all the magnetic moment vectors are aligned in one direction and these substances show magnetic properties even in the absence of external magnetic field.


32. (b): Stronger the acid, more will be the heat of neutralization. Negative sign signifies evolution of heat.



$$= \frac{1}{2}[(B.E.)_{H-H} + (B.E.)_{Cl-Cl}] - (B.E.)_{HCl}$$

$$= \frac{1}{2}[104 + 58] - 103 = -22 \text{ kcal mol}^{-1}$$

34. (b): When the transition metals are in their highest oxidation states, they no longer have tendency to give away electrons, thus they are not basic but show acidic character and form anionic complexes.



36. (a): As the size of the lanthanide ions decreases from  $Ce^{3+}$  to  $Lu^{3+}$ , the covalent character of  $M-OH$  bond increases and hence, the basic strength decreases.

### Your favourite MTG Books/Magazines available in KERALA at


- DHYAN PRAKASHAN BOOK, OPP. VAZHUTHACAUD, TRIVANDRUM PH : 9497430066
- IDEAL BOOK CENTRE, PULIMOODU, TRIVANDRUM PH: 9645163611
- H & C STORE, MAVOOR ROAD, CALICUT PH : 0495-2720620
- H & C STORES-TDM HALL-ERNAKULAM, PH: 0484-2352133/2351233
- VIDYA BOOK HOUSE- KASARAGOD, Mobile : 9447727212
- H & C STORE-KOLLAM, PH : 0474-2765421
- H AND C STORE – KOTTAYAM, PH : 0481-2304351
- TBS PUBLISHERS AND DISTRIBUTORS, KOZHIKODE, PH : 0495-2721025, 2720086, 2721414
- GIFTALIA BOOK BHAVAN-THRISSURE, PH : 0487-2336918

Visit "MTG IN YOUR CITY" on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call **0124-6601200** for further assistance.

Thus,  $\text{Ce}(\text{OH})_3$  is most basic while  $\text{Lu}(\text{OH})_3$  is least basic.


37. (b): Orthoboric acid is a weak monobasic acid with  $K_a = 5.6 \times 10^{-10}$ . It does not act as protonic acid (i.e., proton donor) but behaves as Lewis acid by accepting a pair of electrons from  $\text{OH}^-$  ion.  
 $\text{B}(\text{OH})_3 + 2\text{H}-\text{O}-\text{H} \rightarrow [\text{B}(\text{OH})_4]^- + \text{H}_3\text{O}^+$

38. (c): When bromine water is added to phenol at room temperature, direct bromination gives white precipitate of 2,4,6-tribromophenol.



39. (c): Arsenic drugs (salvarsan) are mainly used in treatment of syphilis.

40. (b):  $\text{K}_2\text{Cr}_2\text{O}_7$  acts as an oxidising agent in presence of dil.  $\text{H}_2\text{SO}_4$ .




Equivalent weight of  $\text{K}_2\text{Cr}_2\text{O}_7$

$$= \frac{\text{Molecular weight}}{\text{Number of electrons gained}} = \frac{M}{6}$$

41. (a): The physical adsorption isobar shows a decrease in  $x/m$  throughout with rise in temperature.

42. (d):  $\text{PhSO}_2\text{Cl} + \text{RR}'\text{NH} \rightarrow \text{PhSO}_2\text{NRR}'$   
*N, N*-dialkyl sulphonamide  
 (Insoluble in  $\text{KOH}$ )

43. (c): All aldehydes except formaldehyde ( $\text{HCHO}$ ) give secondary alcohols with Grignard reagent ( $\text{RMgX}$ ).



44. (c): In case of dilute solutions of acids,  $[\text{H}^+]$  of water cannot be neglected.

$$10^{-8} \text{ M HCl} = 10^{-8} \text{ M H}^+$$

$$\text{Also from } \text{H}_2\text{O}, [\text{H}^+] = 10^{-7} \text{ M}$$

$$\text{Total } [\text{H}^+] = 10^{-7} + 10^{-8} = 10^{-7} (1 + 0.1) = 10^{-7} (1.1)$$




$$\text{Hence, pH} = 7 - 0.0414 = 6.96$$

45. (c): When coordination number is 8,  $r_+/r_-$  ratio lies between 0.732 – 1.

**mtg**

## A Series of Quick Reference Books

Cover complete syllabus in point form • A Handy High Yield Facts Book  
 Easy to Grasp • Essential for all Competitive Examinations



Buy ONLINE at  
[www.mtg.in](http://www.mtg.in)


Buy ONLINE at  
[www.mtg.in](http://www.mtg.in)




**MTG BOOKS**

Plot 99, Sector 44 Institutional Area, Gurgaon (HR) - 122 003  
 Tel.: (0124) 6601200 e-mail : [info@mtg.in](mailto:info@mtg.in) web : [www.mtg.in](http://www.mtg.in)


# The most comprehensive question bank books that you cannot afford to ignore



₹ 275



₹ 275



₹ 300

29 Years' Physics, Chemistry & Biology contain not only chapterwise questions that have appeared over the last 29 years in CBSE's PMT/NEET, but also full solutions, that too by experts. Needless to say, these question banks are essential for any student to compete successfully in NEET.

## HIGHLIGHTS:

- Chapterwise questions of last 29 years' (2016-1988) of CBSE-PMT/NEET
- Chapterwise segregation of questions to help you assess the level of effort required to succeed
- Fully solved questions of NEET (Phase -1) 2016 included
- An unmatched question bank series with close to 1,000 pages having detailed solutions by experts



Scan now with your smartphone or tablet\*

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!



# CONCEPT BOOSTER


## BEWARE OF BASES

Hello students!! When there was a discussion about ACIDS in the last article, how can it be possible that BASES will not come this month!! I am really thankful to all my avid readers. I believe you will like this article same as before. Thanks for the support. Stay healthy.. ☺☺

\*Arunava Sarkar


### BASICITY CORNER

- **Concept 1 :** Basicity is a thermodynamic term. Basicity is better defined with the overall stability of the process during proton abstraction *e.g.*,




Now, if  $\text{BH}^+$  is stable enough, then the overall equilibrium will be shifted towards the right and base  $B$ : is considered to be a good base. If  $\text{BH}^+$  is not well stabilized then the equilibrium will be towards left and  $B$ : is not considered to be a good base.

The stability of  $\text{BH}^+$  is determined by the polarisability of  $\text{BH}^+$  bond. Greater the polarisability, lesser will be the  $\Delta H^\circ$  for bond dissociation (*i.e.* bond dissociation energy) and abstracted  $\text{H}^+$  will be released which will in turn decrease the basicity. The fact is :




Now, if  $\text{A}^-$  is electron donating then it will decrease acidity (antagonistic to the attraction force of oxygen towards the bonding electrons with hydrogen) but if  $\text{A}^-$  is electron withdrawing then it will increase acidity (reinforcing the attraction force of oxygen towards the bonding electrons with hydrogen). So, with the increase in the bond polarisability of  $\text{B}-\text{H}$  bond, basicity decreases.


As per this, the basicity order of period 2 elements is :



For the similar reason, along the group, polarisability increases and basicity decreases, so,  $\text{R}\ddot{\text{O}} > \text{R}\ddot{\text{S}}:$ . As per the electron donating or electron withdrawing tendency of the attached group(s), basicity alters, *e.g.*,



- **Concept 2 :** In general, with increase in the number of alkyl groups attached with electron donating atom, basicity increases, *e.g.*,



But a few exceptions are there which are given in the following table :


**Table 1 :**

Compound	$\text{p}K_b$	
$:\text{NH}_3$	4.5	
$\text{CH}_3\ddot{\text{N}}\text{H}_2$	3.36	
$\text{CH}_3\text{>}\ddot{\text{N}}\text{H}$	3.23	surprising
$\text{H}_3\text{C}>\ddot{\text{N}}:$	4.2	


\* Institute of Chemistry (IOC)- Asansol, Durgapur, Dhanbad, Burdwan, Kolkata, Jamshedpur, Bokaro, Patna



# BEST TOOLS FOR SUCCESS IN JEE Main



₹ 325



₹ 375

**10** Very Similar Practice Test Papers

**15** Years JEE MAIN 2016-2015 (Offline & Online)-2013  
& AIEEE (2012-2002)



Available at all leading book shops throughout the country.  
For more information or for help in placing your order:  
Call 0124-6601200 or email: [info@mtg.in](mailto:info@mtg.in)


Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!

**Table 2 :**

Compound	$pK_b$	
$\text{Et}\ddot{\text{N}}\text{H}_2$	3.33	
$\begin{array}{c} \text{Et} \\ \diagup \\ \text{N} \ddot{\text{H}} \\ \diagdown \\ \text{Et} \end{array}$	3.07	surprising
$\begin{array}{c} \text{Et} \\ \diagup \\ \text{N} \\ \diagdown \\ \text{Et} \end{array}$	3.12	


This only happens in water or a solvent with proton ( $\text{H}^+$ ). But in gaseous phase normal order *i.e.*,  $3^\circ > 2^\circ > 1^\circ$  amine is observed.

○ **Concept 3 :**  $3^\circ > 2^\circ > 1^\circ$




$\text{R}_4\text{N}^+\ddot{\text{O}}\text{H}^-$  is reported to be completely ionized and it has basicity comparable with mineral alkalies.


○ **Concept 4 :** Guanidine is an extremely strong base with  $pK_b = 0.4$ . Guanidine has the following structure :




and this is expected to show the following resonating structures :



When guanidine accepts a proton (*i.e.* it acts as a base),



The conjugate acid formed has the following resonating structures :



Now, a few points which are to be considered are :

1. Structures II and III are having charge separation and this is not good as per the

overall stability of the resonating structures is considered whereas in structures IV, V and VI there is no charge separation, only the +ve charge is distributed here which is a good sign as per the stability of resonating structures is considered. So, I is not as resonance stabilised as IV.

2. I is not equivalent with II and III. IV is equivalent with V and VI. Equivalent structures give better stability.
3. For protonation of guanidine, a very small amount of energy is required. So, energy difference is not of much importance.

Guanidine is definitely a strong base. Structures are said to be equivalent on the basis of the energy content and symmetry.

Guanidine has  $pK_b$  value of 0.4 but a similar structure of amidine *i.e.*,  $\text{RC}-\text{NH}_2$  has  $pK_b = 1.6$



with  $\text{R} = -\text{CH}_3$



This is due to the lesser stability of the corresponding conjugate acid of amidine in comparison with the conjugate acid of guanidine.

**Your favourite MTG Books/Magazines available in ODISHA at**


- **SRI RAM BOOKS-BHUBANESWAR,**  
Ph: 0674-2391385
- **KBRC BOOK STALL-BHUBANESWAR,**  
Ph: 9937006506
- **PADMALAYA BOOK SELLER-BHUBANESWAR**  
Ph: 0674-2396922, 3039692
- **PRAGNYA-BHUBANESWAR,**  
Ph: 0674-2395757, 30396922
- **SRI MADHAB BOOK STORE-CUTTACK,**  
Ph: 0671-2431158, 2431148
- **KITAB MAHAL – CUTTACK,**  
Ph: 0671-2648333, 2648301
- **A.K. MISHRA AGENCY PVT. LTD.-CUTTACK,**  
Ph: 0671-2322244, 2322255, 2322266, 2332233

Visit **"MTG IN YOUR CITY"** on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call **0124-6601200** for further assistance.

Suppose, the following base is given :



○ **Concept 5 :** Which one is more basic?



As imidazoline contains two nitrogen atoms whereas pyridine contains only one hence, imidazoline is a better base. The following table confirms it :


**Table 3 :**

Compound	$pK_b$
Imidazoline	3
Pyridine	7.8


○ **Concept 6 :** Aromatic amine *i.e.*, aniline is less basic than ammonia because the lone pair over nitrogen

atom takes part in resonance with the benzene ring. This can be understood from the  $pK_b$  values given in the following table :

**Table 4 :**

Compound	$pK_b$	
$\text{NH}_3$	4.75	 basicity decreases a lot
$\text{C}_6\text{H}_5\text{NH}_2$	9.38	
$\text{C}_6\text{H}_5\text{NHMe}$	9.16	
$\text{C}_6\text{H}_5\text{NMe}_2$	8.85	

In case of  $\text{C}_6\text{H}_5\text{NMe}_2$ , there is a steric repulsion between the *ortho* hydrogen and the bulky  $-\text{NMe}_2$  group.



Due to this steric repulsion, lone pair of nitrogen can show basicity.

## SOLUTIONS OF MAY 2016 CROSSWORD


[illegible]

## Winners of April 2016 Crossword


- Hrishikesh Sarmah, Assam
- Samrat Gupta, Kolkata



The most Reliable and Featured  
**22 Years' AIIMS EXPLORER and**  
**AIIMS CHAPTERWISE SOLUTIONS** in the market



₹ 400



₹ 375

#### HIGHLIGHTS:

- 22 years' (1994-2015) Solved Papers with Detailed Solutions
- 10 Model Test Papers
- 600+ General Knowledge Questions
- 1600+ Assertion Reason Questions
- 22 years' (1994-2015) Chapterwise Solutions
- Subjectwise distribution of 22 years' questions


# NCERT CORNER

The questions given in this column have been prepared strictly on the basis of NCERT Chemistry. Papers of JEE(Main & Advanced)/NEET/AIIMS/other PMTs are drawn heavily from NCERT books. Practice Hard! All the best!!




- 1 g of Mg is burnt in a closed vessel containing 0.5 g of O<sub>2</sub>. Which of the following statements is correct?
  - O<sub>2</sub> is the limiting reagent and Mg is in excess by 0.25 g.
  - Mg is the limiting reagent and is in excess by 0.5 g.
  - O<sub>2</sub> is the limiting reagent and is in excess by 0.25 g.
  - O<sub>2</sub> is the limiting reagent and Mg is in excess by 0.75 g.
- Hydrogen bonds are formed in many compounds like H<sub>2</sub>O, HF and NH<sub>3</sub>. The correct decreasing order of boiling points of these compounds is
  - HF > H<sub>2</sub>O > NH<sub>3</sub>
  - H<sub>2</sub>O > HF > NH<sub>3</sub>
  - NH<sub>3</sub> > HF > H<sub>2</sub>O
  - NH<sub>3</sub> > H<sub>2</sub>O > HF
- Brine is electrolysed by using inert electrodes. The reaction at anode is
  - $\text{Cl}^-_{(aq)} \longrightarrow \frac{1}{2} \text{Cl}_{2(g)} + e^-$ ;  $E^\circ_{\text{cell}} = 1.36 \text{ V}$
  - $2\text{H}_2\text{O}_{(l)} \longrightarrow \text{O}_{2(g)} + 4\text{H}^+ + 4e^-$ ;  $E^\circ_{\text{cell}} = 1.23 \text{ V}$
  - $\text{Na}^+_{(aq)} + e^- \longrightarrow \text{Na}_{(s)}$ ;  $E^\circ_{\text{cell}} = -2.71 \text{ V}$
  - $\text{H}^+_{(aq)} + e^- \longrightarrow \frac{1}{2} \text{H}_{2(g)}$ ;  $E^\circ_{\text{cell}} = 0.00 \text{ V}$
- An organic compound 'A' having molecular formula C<sub>3</sub>H<sub>6</sub> on treatment with aqueous H<sub>2</sub>SO<sub>4</sub> gives 'B' which on treatment with HCl/ZnCl<sub>2</sub> gives 'C'. The compound 'C' on treatment with ethanolic KOH gives back the compound 'A'. Identify the compound 'C'.
  - CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>Cl
  - CH<sub>3</sub>—CH=CH<sub>2</sub>
  - CH<sub>3</sub>CH(Cl)CH<sub>3</sub>
  - CH<sub>3</sub>CH<sub>2</sub>CH<sub>3</sub>
- Considering the elements F, Cl, O and N, the correct order of their chemical reactivity in terms of oxidising property is
  - F > Cl > O > N
  - F > O > Cl > N
  - Cl > F > O > N
  - O > F > N > Cl
- The pollutants which come directly in the air from sources are called primary pollutants. Primary pollutants are sometimes converted into secondary pollutants. Which of the following belongs to secondary air pollutants?
  - CO
  - Hydrocarbon
  - Peroxyacetyl nitrate
  - NO
- Which of the following is homopolymer?
  - PMMA
  - Bakelite
  - Glyptal
  - Nylon-6,6
- Which of the following reactions increases the production of dihydrogen from synthesis gas?
  - $\text{CH}_4(g) + \text{H}_2\text{O}(g) \xrightarrow[\text{Ni}]{1270 \text{ K}} \text{CO}(g) + 3\text{H}_2(g)$
  - $\text{C}(s) + \text{H}_2\text{O}(g) \xrightarrow{1270 \text{ K}} \text{CO}(g) + \text{H}_2(g)$
  - $\text{CO}(g) + \text{H}_2\text{O}(g) \xrightarrow[\text{Catalyst}]{673 \text{ K}} \text{CO}_2(g) + \text{H}_2(g)$
  - $\text{C}_2\text{H}_6(g) + 2\text{H}_2\text{O}(g) \xrightarrow[\text{Ni}]{1270 \text{ K}} 2\text{CO}(g) + 5\text{H}_2(g)$
- If  $q_1$  is the specific charge ( $e/m$ ) of cathode rays and  $q_2$  is that of positive rays then
  - $q_1 = q_2$
  - $q_1 < q_2$
  - $q_1 > q_2$
  - none of these.
- Alum is added to water containing suspended impurities
  - to make a colloidal solution
  - to coagulate the suspended impurities
  - to remove impurities of calcium and magnesium
  - to protect the colloidal solution from getting precipitated.
- Which of the following reactions will not yield phenol?
  -













12. How many years would it take to spend Avogadro number of rupees at the rate of 10 lac rupees per second?

- (a)  $1.90 \times 10^{10}$  years (b)  $6.02 \times 10^3$  years  
(c)  $1.90 \times 10^6$  years (d)  $4.01 \times 10^4$  years

13. Which product(s) is/are formed when the compound -CHO is treated with concentrated aqueous KOH solution?

- (a) -CHO
- (b)  and 
- (c)  and 
- (d)  and 

14. Predict the structure of the product in the given reaction.



15. Which is the correct decreasing order of reactivity towards  $\text{S}_{\text{N}}2$  displacement of the following compounds?

- (a) 1-Bromo-2-methylbutane > 1-Bromo-2,2-dimethylpropane > 1-Bromo-3-methylbutane > 1-Bromobutane  
(b) 1-Bromo-2,2-dimethylpropane > 1-Bromobutane > 1-Bromo-2-methylbutane > 1-Bromo-3-methylbutane  
(c) 1-Bromobutane > 1-Bromo-2-methylbutane > 1-Bromo-2,2-dimethylpropane > 1-Bromo-3-methylbutane  
(d) 1-Bromobutane > 1-Bromo-3-methylbutane > 1-Bromo-2-methylbutane > 1-Bromo-2,2-dimethylpropane

16. Corrosion of iron is essentially an electrochemical phenomenon where the cell reactions are

- (a) Fe is oxidised to  $\text{Fe}^{2+}$  and dissolved oxygen in water is reduced to  $\text{OH}^-$   
(b) Fe is oxidised to  $\text{Fe}^{3+}$  and  $\text{H}_2\text{O}$  is reduced to  $\text{O}_2^{2-}$   
(c) Fe is oxidised to  $\text{Fe}^{2+}$  and  $\text{H}_2\text{O}$  is reduced to  $\text{O}_2^-$   
(d) Fe is oxidised to  $\text{Fe}^{2+}$  and  $\text{H}_2\text{O}$  is reduced to  $\text{O}_2$ .

17. For the given reactions (i) and (ii) which of the algebraic relations is correct?

- (i)  $\text{C}_{(\text{g})} + 4\text{H}_{(\text{g})} \rightarrow \text{CH}_{4(\text{g})}$ ;  $\Delta_r H = x \text{ kJ mol}^{-1}$   
(ii)  $\text{C}_{(\text{s})} (\text{graphite}) + 2\text{H}_{2(\text{g})} \rightarrow \text{CH}_{4(\text{g})}$ ;  $\Delta_r H = y \text{ kJ mol}^{-1}$

- (a)  $x = y$  (b)  $x = 2y$  (c)  $x > y$  (d)  $x < y$

18. Which of the following is a broad spectrum antibiotic?

- (a) Ampicillin (b) Penicillin  
(c) Chloramphenicol (d) Terpineol


19. When a substance 'A' reacts with water it produces a combustible gas 'B' and a solution of substance 'C' in water. When another substance 'D' reacts with this solution of 'C', it also produces the same gas 'B'. 'A' imparts a deep golden yellow colour to smokeless flame of Bunsen burner. A, B, C and D are respectively

- (a) Na,  $\text{H}_2$ , NaOH and Zn  
(b) K,  $\text{H}_2$ , KOH and Al  
(c) Ca,  $\text{H}_2$ ,  $\text{Ca}(\text{OH})_2$  and Sn  
(d)  $\text{CaC}_2$ ,  $\text{C}_2\text{H}_2$ ,  $\text{Ca}(\text{OH})_2$  and Fe

20. Identify the correct order of acidic strengths of  $\text{CO}_2$ , CuO, CaO and  $\text{H}_2\text{O}$ .

- (a)  $\text{CaO} < \text{CuO} < \text{H}_2\text{O} < \text{CO}_2$   
(b)  $\text{H}_2\text{O} < \text{CuO} < \text{CaO} < \text{CO}_2$   
(c)  $\text{CaO} < \text{H}_2\text{O} < \text{CuO} < \text{CO}_2$   
(d)  $\text{H}_2\text{O} < \text{CO}_2 < \text{CaO} < \text{CuO}$


21. Consider a first order gas phase decomposition reaction given below :



The initial pressure of the system before decomposition of A was  $P_i$ . After lapse of time ' $t$ ', total pressure of the system increased by  $x$  units and became ' $P_t$ '. The rate constant  $k$  for the reaction is given as

- (a)  $k = \frac{2.303}{t} \log \frac{P_i}{P_i - x}$   
 (b)  $k = \frac{2.303}{t} \log \frac{P_i}{2P_i - P_t}$   
 (c)  $k = \frac{2.303}{t} \log \frac{P_i}{2P_i + P_t}$   
 (d)  $k = \frac{2.303}{t} \log \frac{P_i}{P_i + x}$

22. Which of the following represents the product formed in the given reaction?




- (a) (b)   
 (c) (d)

23. The critical temperature of water is higher than that of  $O_2$  because the  $H_2O$  molecule has

- (a) fewer electrons than  $O_2$   
 (b) two covalent bonds  
 (c) V-shape  
 (d) dipole moment.


24. The standard electrode potentials of few metals are given below :




Arrange these metals in increasing order of their reducing power.

- (a)  $Hg < Cr < Ag < Mg < K$   
 (b)  $Hg < K < Mg < Cr < Ag$   
 (c)  $Ag < K < Mg < Hg < Cr$   
 (d)  $Ag < Hg < Cr < Mg < K$

25. For the equilibrium,



which of the given plots should be useful to determine the enthalpy change of the reaction ( $\Delta H$ )?



26. When aqueous solution of potassium fluoride is added to the blue coloured aqueous  $CuSO_4$  solution, a green precipitate is formed. This is because

- (a) on adding KF,  $H_2O$  being weak field ligand is replaced by  $F^-$  ions forming  $[CuF_4]^{2-}$  which is green in colour  
 (b) potassium is coordinated to  $[Cu(H_2O)_4]^{2+}$  ion present in  $CuSO_4$  and gives green colour  
 (c) on adding KF,  $Cu^{2+}$  are replaced by  $K^+$  forming a green complex  
 (d) blue colour of  $CuSO_4$  and yellow colour of KI form green colour on mixing.

27. Which of the following statements is incorrect?

- (a) In equilibrium mixture of ice and water kept in perfectly insulated flask, mass of ice and water does not change with time.  
 (b) The intensity of red colour increases when oxalic acid is added to a solution containing iron (III) nitrate and potassium thiocyanate.  
 (c) On addition of catalyst the equilibrium constant value is not affected.  
 (d) Equilibrium constant for a reaction with negative  $\Delta H$  value decreases as the temperature increases.

28.  $P_4O_6$  reacts with water according to the equation  $P_4O_6 + 6H_2O \rightarrow 4H_3PO_3$ . Calculate the volume of 0.1 M NaOH solution required to neutralise the acid formed by dissolving 1.1 g of  $P_4O_6$  in  $H_2O$ .

- (a) 0.004 L (b) 0.4 L  
 (c) 4.4 L (d) 1.2 L

29. Negative electronic energy (negative sign for all values of energy) for hydrogen atom means

- (a) the energy of an electron in the atom is lower than the energy of a free electron at rest which is taken as zero  
 (b) when the electron is free from the influence of nucleus it has a negative value which becomes more negative

- (c) when the electron is attracted by the nucleus the energy is absorbed which means a negative value  
 (d) energy is released by hydrogen atom in ground state.
30. White metallic tin gets converted to grey powder at low temperature. This transformation is related to  
 (a) an interaction with nitrogen of the air at very low temperatures  
 (b) a change in the crystalline structure of tin  
 (c) an interaction with water vapour contained in the humid air  
 (d) a change in the partial pressure of oxygen in the air.
31. In the Lassaigne's test for nitrogen in an organic compound, the prussian blue colour is obtained due to the formation of  
 (a)  $\text{Na}_4[\text{Fe}(\text{CN})_6]$  (b)  $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$   
 (c)  $\text{Fe}_2[\text{Fe}(\text{CN})_6]$  (d)  $\text{Fe}_3[\text{Fe}(\text{CN})_6]_4$
32. Magnetic moment of  $\text{Cr}^{2+}$  is nearest to  
 (a)  $\text{Fe}^{2+}$  (b)  $\text{Mn}^{2+}$  (c)  $\text{Co}^{2+}$  (d)  $\text{Ni}^{2+}$
33. A graph of adsorption isobar of chemisorption shows that adsorption  
 (a) first decreases with temperature and then increases  
 (b) first increases with temperature and then decreases  
 (c) increases with temperature  
 (d) decreases with temperature.
34. An aqueous solution contains equimolar quantities of  $\text{Zn}(\text{NO}_3)_2$ ,  $\text{AgNO}_3$ ,  $\text{Cu}(\text{NO}_3)_2$ .  
 $E^\circ$  values of  
 $\text{Zn}^{2+}/\text{Zn} = -0.76 \text{ V}$ ;  $\text{Ag}^+/\text{Ag} = +0.80 \text{ V}$ ,  
 $\text{Cu}^{2+}/\text{Cu} = +0.34$ .  
 As the voltage is increased, the sequence in which metals are expected to be deposited is  
 (a) Cu, Ag (b) Ag, Cu  
 (c) Cu, Zn (d) Zn, Cu, Ag
35. Most common types of secondary structures of proteins are  
 (a)  $\alpha$ -helix and  $\beta$ -helix structures  
 (b)  $\alpha$ -helix and  $\beta$ -pleated sheet structures  
 (c) right and left hand twisted structures  
 (d) globular and fibrous structures.
36. For a gaseous reaction  
 $2\text{A} + \text{B}_2 \longrightarrow 2\text{AB}$ ,


The following rate data were obtained at 300 K.

Expt.	Concentration		Rate of disappearance of $\text{B}_2$ ( $\text{mol L}^{-1} \text{ min}^{-1}$ )
	[A]	[ $\text{B}_2$ ]	
1.	0.015	0.15	$1.8 \times 10^{-3}$
2.	0.09	0.15	$1.08 \times 10^{-2}$
3.	0.015	0.45	$5.4 \times 10^{-3}$

What is the rate law?

- (a)  $r = k[\text{A}][\text{B}_2]$  (b)  $r = [\text{A}]^2 [\text{B}_2]^1$   
 (c)  $r = k[\text{A}][\text{B}_2]^2$  (d)  $r = k[\text{B}_2]$
37. The drain cleaner, Drainex contains small bits of aluminium which react with caustic soda to produce dihydrogen. What volume of dihydrogen at  $20^\circ\text{C}$  and one bar will be released when 0.15 g of aluminium reacts?  
 (a) 204 mL (b) 200 mL  
 (c) 203 mL (d) 400 mL
38. The molecular shapes of  $\text{SF}_4$ ,  $\text{CF}_4$  and  $\text{XeF}_4$  are  
 (a) the same with 2, 0 and 1 lone pairs of electrons on the central atom, respectively  
 (b) the same with 1, 1 and 1 lone pair of electrons on the central atom, respectively  
 (c) different with 0, 1 and 2 lone pairs of electrons on the central atom, respectively  
 (d) different with 1, 0 and 2 lone pairs of electrons on the central atom, respectively.
39. Which of the following is a correct statement?  
 (a)  $(\text{SiH}_3)_3\text{N}$  is a stronger base than  $(\text{CH}_3)_3\text{N}$ .  
 (b)  $(\text{CH}_3)_3\text{N}$  has a pyramidal structure.  
 (c)  $(\text{SiH}_3)_3\text{N}$  has a pyramidal structure.  
 (d)  $\text{SiO}_4^{4-}$  units are not tetrahedral.
40. The addition of HBr to 1-butene gives a mixture of products (I), (II) and (III).  
 (I)  $\text{H}_5\text{C}_2 - \overset{\text{Br}}{\underset{\text{H}}{\text{C}}} - \text{CH}_3$  (II)  $\text{H} - \overset{\text{C}_2\text{H}_5}{\underset{\text{Br}}{\text{C}}} - \text{CH}_3$   
 (III)  $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{Br}$   
 The mixture consists of  
 (a) (I) and (II) as major and (III) as minor products  
 (b) (II) as major, (I) and (III) as minor products  
 (c) (II) as minor, (I) and (III) as major products  
 (d) (I) and (II) as minor and (III) as major products.


41. Of the following reaction,



X, Y and Z are respectively

- (a) sodalime,  $\text{NaOH}_{(aq)}$ ,  $\text{CH}_3\text{Cl}$   
 (b)  $\text{NaOH}_{(aq)}$ , sodalime,  $\text{CH}_3\text{Cl}$   
 (c)  $\text{CH}_3\text{Cl}$ ,  $\text{NaOH}_{(aq)}$ , sodalime  
 (d) sodalime,  $\text{CH}_3\text{Cl}$ ,  $\text{NaOH}_{(aq)}$ .
42. The edge length of unit cell of a metal having molecular weight 75 g/mol is  $5 \text{ \AA}$  which crystallizes in body centred cubic lattice. If the density is 2 g/cc then find the radius in pm of metal atom. ( $N_A = 6 \times 10^{23}$ ).  
 (a) 116.5 pm (b) 316.5 pm  
 (c) 216.5 pm (d) 416.5 pm
43. X-ray diffraction studies show that copper crystallises in fcc unit cell with cell edge of  $3.608 \times 10^{-8} \text{ cm}$ . In a separate experiment, copper is determined to have a density of  $8.92 \text{ g/cm}^3$ . The atomic mass of copper is  
 (a) 160 u (b) 35 u  
 (c) 63 u (d) 25 u

44. In the reaction,



What is not true about the  $-\text{NH}-\text{C}(=\text{O})-\text{CH}_3$  group?

- (a) It reduces the activating influence of  $-\text{NH}_2$  group.  
 (b) The group is *ortho-para* directing.  
 (c) The group can be converted into  $-\text{NH}_2$  by adding a dilute acid.  
 (d) When this group is present on the benzene ring, it becomes very easy to carry on nucleophilic substitution.
45. In Duma's method 0.52 g of an organic compound on combustion gave 68.6 mL  $\text{N}_2$  at  $27^\circ\text{C}$  and 756 mm pressure. What is the percentage of nitrogen in the compound?  
 (a) 12.22% (b) 14.93%  
 (c) 15.84% (d) 16.23%

46. Amongst the given set of reactants, the most appropriate for preparing  $2^\circ$  amine is

- (a)  $2^\circ \text{R}-\text{Br} + \text{NH}_3$   
 (b)  $2^\circ \text{R}-\text{Br} + \text{NaCN}$  followed by  $\text{H}_2/\text{Pt}$   
 (c)  $1^\circ \text{R}-\text{NH}_2 + \text{RCHO}$  followed by  $\text{H}_2/\text{Pt}$   
 (d)  $1^\circ \text{R}-\text{Br}$  (2 mol) + potassium phthalimide followed by  $\text{H}_3\text{O}^+/\text{heat}$ .

47. Which of the following statements is not correct?


- (a) 5% aqueous solutions of  $\text{NaCl}$  and  $\text{KCl}$  are said to be isomolar.  
 (b) 1 M sucrose solution and 1 M glucose solution are isotonic.  
 (c) Molecular mass of acetic acid or benzoic acid is higher than normal mass in cryoscopic methods.

(d) For the same solution,  $\frac{\Delta T_b}{\Delta T_f} = \frac{K_b}{K_f}$ .


48. The correct order of stability is

- (a)  $\text{SnF}_2 > \text{PbF}_2 > \text{GeF}_2$   
 (b)  $\text{GeF}_2 > \text{SnF}_2 > \text{PbF}_2$   
 (c)  $\text{GeF}_2 > \text{PbF}_2 > \text{SnF}_2$   
 (d)  $\text{PbF}_2 > \text{SnF}_2 > \text{GeF}_2$

49. The correct energy level diagram for  $[\text{Co}(\text{CN})_6]^{3-}$  is




50. 1.0 mol of a monoatomic ideal gas is expanded from state (1) to (2) as shown in figure. Calculate the work done for the expansion of gas from state (1) to (2) at 298 K.
- (a) - 1717.63 J                      (b) - 5705.8 J  
(c) + 1717.63 J                      (d) + 5705.8 J




### SOLUTIONS

1. (a):  $2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$   
 $2 \times 24 \quad 2 \times 16 \quad 2(24 + 16)$   
 48 g of Mg requires 32 g of  $\text{O}_2$   
 1 g of Mg requires  $\frac{32}{48} = 0.66$  g of  $\text{O}_2$   
 Oxygen available = 0.5 g  
 Hence,  $\text{O}_2$  is the limiting reagent.  
 32 g of  $\text{O}_2$  reacts with 48 g of Mg  
 0.5 g of  $\text{O}_2$  will react with  $\frac{48}{32} \times 0.5 = 0.75$  g of Mg  
 Excess of Mg =  $1.0 - 0.75 = 0.25$  g
2. (b): Strength of H-bond is in the order :  
 $\text{H} \cdots \text{F} > \text{H} \cdots \text{O} > \text{H} \cdots \text{N}$   
 But each  $\text{H}_2\text{O}$  molecule is linked to 4 other  $\text{H}_2\text{O}$  molecules through H-bonds whereas each HF molecule is linked only to two other HF molecules.  
 Hence, boiling point of  $\text{H}_2\text{O} >$  boiling point of HF.
3. (a): For oxidation at anode, two possible reactions are oxidation of chlorine and of oxygen. Out of these two, oxidation of chlorine ion is preferred because oxidation of oxygen requires overvoltage.  
 Chlorine is obtained by electrolysis giving out hydrogen and aqueous NaOH as by-products.

4. (c):




5. (b): Within a period, the oxidising character increases from left to right. Therefore, among F, O and N, oxidising power decreases in the order :  $\text{F} > \text{O} > \text{N}$ . However, within a group, oxidising power decreases from top to bottom. Thus, F is a stronger oxidising agent than Cl. Further because O is more electronegative than Cl, therefore, O is a stronger oxidising agent than Cl. Thus, overall decreasing order of oxidising power is :  $\text{F} > \text{O} > \text{Cl} > \text{N}$ .
6. (c): Peroxyacetyl nitrate (PAN) formed from the primary pollutants  $\text{NO}_2$ ,  $\text{O}_3$  and hydrocarbons is the secondary pollutant.
7. (a): PMMA is homopolymer *i.e.*, polymethyl-methacrylate.




The production of dihydrogen can be increased by reacting CO of syngas mixture with steam in presence of iron chromate as a catalyst. This is called water-gas shift reaction.

9. (c): Mass of positively charged ions in positive rays is more than the mass of electrons.
10. (b): Water obtained from natural sources often contains suspended impurities. Alum is added to such water to coagulate the suspended impurities and make water fit for drinking purposes.
11. (d): Chlorobenzene does not undergo hydrolysis on treatment with aqueous NaOH at 298 K.
12. (a): Avogadro number =  $6.023 \times 10^{23}$   
 Total rupees =  $6.023 \times 10^{23}$   
 Rate of spending = 10 lac rupees/sec =  $10^6$  rupees/sec  
 Total amount spent in one year  
 $= 10^6 \times 60 \times 60 \times 24 \times 365$  rupees  
 $\therefore$  Number of years to spend all the rupees  
 $= \frac{6.023 \times 10^{23} \text{ rupees}}{10^6 \times 60 \times 60 \times 24 \times 365 \text{ rupees/year}}$   
 $= 1.90988 \times 10^{10}$  years
13. (b): Benzaldehyde when treated with conc. KOH solution undergoes Cannizzaro reaction. One molecule of aldehyde is reduced and other is oxidised.








14. (c) :



15. (d) :  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$   
1-Bromobutane  
(1° with no branching)



Since in case of 1° alkyl halides, steric hindrance increases in the order : *n*-alkyl halides, alkyl halide with a substituent at any position other than the β-position, one substituent at the β-position, two substituents at the β-position, therefore, the reactivity decreases in the same order. Thus, the reactivity of the given alkyl bromides decreases in the order :


1-Bromobutane > 1-Bromo-3-methylbutane >

1-Bromo-2-methylbutane > 1-Bromo-2,2-dimethylpropane

16. (a) : At anode :  $\text{Fe} \longrightarrow \text{Fe}^{2+} + 2e^-$

At cathode :  $\text{O}_2 + 2\text{H}_2\text{O} + 4e^- \longrightarrow 4\text{OH}^-$


The overall reaction is



$\text{Fe}(\text{OH})_2$  may be dehydrated to iron oxide  $\text{FeO}$ , or further oxidised to  $\text{Fe}(\text{OH})_3$  and then dehydrated to iron rust,  $\text{Fe}_2\text{O}_3$ .

17. (c) :  $x > y$  because same bonds are formed in reactions (i) and (ii) but bonds between reactant molecules are broken only in reaction (ii). As energy is absorbed when bonds are broken, energy released in reaction (i) is greater than that in reaction (ii).


18. (c) : Chloramphenicol is a broad spectrum drug and can be used in treatment of typhoid, dysentery, meningitis, etc.



Na (A) imparts golden yellow colour to the flame.

20. (a) :  $\text{CaO}$  is basic while  $\text{CO}_2$  is most acidic of these. The increasing acidic strength order is  $\text{CaO} < \text{CuO} < \text{H}_2\text{O} < \text{CO}_2$ .


21. (b) : Consider a first order gas phase decomposition reaction :



The initial pressure of the system before decomposition of A was  $P_i$ .

After lapse of time ( $t$ ), total pressure of the system increased by  $x$  units and became ' $P_t$ '.

In other words, the pressure of A decreased by  $x$ .




$$P_t = (P_i - x) + x + x = P_i + x$$

$$x = P_t - P_i$$

Pressure of A after time  $t$ ,  $P_A = P_i - x$

$$= P_i - P_t + P_i = 2P_i - P_t$$

$$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]} = \frac{2.303}{t} \log \frac{P_i}{2P_i - P_t}$$



23. (d) : More the dipole moment more is the critical temperature.

24. (d) : Lower the electrode potential, better is the reducing power. Since the electrode potentials increase in the order :  $\text{K}^+/\text{K} (-2.93 \text{ V})$ ,  $\text{Mg}^{2+}/\text{Mg} (-2.37 \text{ V})$ ,  $\text{Cr}^{3+}/\text{Cr} (-0.74 \text{ V})$ ,  $\text{Hg}^{2+}/\text{Hg} (0.79 \text{ V})$ ,  $\text{Ag}^+/\text{Ag} (0.80 \text{ V})$ , therefore, reducing power of metals increases in the order :




25. (a) : In this case,  $K_p = p_{\text{H}_2\text{O}}$ ; we know


$$\ln K_p = \ln A - \frac{\Delta H}{RT}$$

∴ Slope will give the value of  $\Delta H$ .


26. (a) : Aqueous  $\text{CuSO}_4$  solution contains  $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$  ions which are blue in colour. When aqueous solution of KF is added,  $\text{H}_2\text{O}$  being weak field ligand can be replaced by  $\text{F}^-$  ions forming  $[\text{CuF}_4]^{2-}$  which is green in colour.



27. (b): When oxalic acid is added to a solution containing iron nitrate and potassium thiocyanate, oxalic acid reacts with  $\text{Fe}^{3+}$  ions to form a stable complex ion  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$  thus, decreasing the concentration of free  $\text{Fe}^{3+}$  ions which in turn decreases the intensity of red colour.



28. (b):  $\text{P}_4\text{O}_6 + 6\text{H}_2\text{O} \rightarrow 4\text{H}_3\text{PO}_3$   
 $[\text{H}_3\text{PO}_3 + 2\text{NaOH} \rightarrow \text{Na}_2\text{HPO}_3 + 2\text{H}_2\text{O}] \times 4$   
 (Neutralisation reaction)



Product formed by 1 mol of  $\text{P}_4\text{O}_6$  is neutralised by 8 moles of NaOH.

$\therefore$  Product formed by  $\frac{1.1}{220}$  mol of  $\text{P}_4\text{O}_6$  will be

neutralised by  $\frac{1.1}{220} \times 8$  moles of NaOH.

Molarity of NaOH solution is 0.1 M.


$\Rightarrow$  0.1 mol NaOH is present in 1 L solution.

$\therefore \frac{1.1}{220} \times 8$  moles NaOH will be present in

$$\frac{1.1 \times 8}{220 \times 0.1} = \frac{88}{220} = \frac{4}{10} = 0.4 \text{ L}$$

29. (a): The negative sign means that the energy of an electron in an atom is lower than the energy of a free electron at rest. When electron is free from the influence of nucleus, the energy is taken as zero. When the electron is attracted by the nucleus, the energy is emitted and lowered.

30. (b): As temperature decreases, white tin ( $\beta$ -form) changes to grey tin ( $\alpha$ -form).



During conversion of white tin to grey tin volume increases. Grey tin is very brittle and easily crumbles down to powder.

31. (b): The prussian blue colour in the Lassaigne's test is due to the formation of ferric ferrocyanide,  $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$ .

32. (a):  $\text{Cr}^{2+} = 3d^4$ , No. of unpaired electrons ( $n$ ) = 4

$$\begin{aligned} \text{Magnetic moment} &= \sqrt{n(n+2)} \text{ BM} \\ &= \sqrt{4(4+2)} = \sqrt{24} = 4.89 \text{ BM} \end{aligned}$$

$\text{Fe}^{2+} = 3d^6$ , No. of unpaired electrons ( $n$ ) = 4

$$\begin{aligned} \text{Magnetic moment} &= \sqrt{4(4+2)} \text{ BM} \\ &= \sqrt{24} = 4.89 \text{ BM} \end{aligned}$$

$\text{Mn}^{2+} = 3d^5$ , No. of unpaired electrons ( $n$ ) = 5

$$\begin{aligned} \text{Magnetic moment} &= \sqrt{5(5+2)} \text{ BM} \\ &= \sqrt{35} = 5.91 \text{ BM} \end{aligned}$$

$\text{Co}^{2+} = 3d^7$ , No. of unpaired electrons ( $n$ ) = 3

$$\begin{aligned} \text{Magnetic moment} &= \sqrt{3(3+2)} \text{ BM} \\ &= \sqrt{15} = 3.87 \text{ BM} \end{aligned}$$

$\text{Ni}^{2+} = 3d^8$ , No. of unpaired electrons ( $n$ ) = 2

$$\begin{aligned} \text{Magnetic moment} &= \sqrt{2(2+2)} \text{ BM} \\ &= \sqrt{8} = 2.82 \text{ BM} \end{aligned}$$


33. (b): A graph of adsorption isobar of chemisorption shows that the adsorption first increases with temperature and then decreases. The initial increase is due to heat supplied, which acts as activation energy required in chemisorption. Later, it decreases due to the exothermic nature of adsorption at equilibrium.

34. (b): Higher the reduction potential of a metal more easily it is deposited at cathode. In an aqueous solution Zn will not be deposited.

35. (b): Due to regular folding of backbone of the peptide chain, two most common secondary structures of proteins are  $\alpha$ -helix and  $\beta$ -pleated sheet structures. In  $\alpha$ -helix, all H-bonds are formed by twisting the chain into a right handed helix. In  $\beta$ -pleated, all peptide chains are stretched to maximum extension and then laid side by side and connected by hydrogen bonds.

36. (a): From expt. (1) and (2), it is clear that when  $[B_2]$  is kept constant and  $[A]$  is made 6 times, the rate also becomes 6 times, thus  $r \propto [A]^1$ . Further from expt (1) and (3) when  $[A]$  is kept constant and  $[B_2]$  is tripled, rate also becomes three times, Thus,  $r \propto [B_2]^1$   
 Hence,  $r = k [A]^1 [B_2]^1$

37. (c): The reaction between aluminium and caustic soda is



$$2 \times 27 = 54 \text{ g} \qquad \qquad \qquad 3 \times 22.4 \text{ L}$$

$\therefore$  54 g of Al produces  $3 \times 22.4$  L of  $\text{H}_2$  at STP

0.15 g of Al will produce  $\frac{3 \times 22.4}{54} \times 0.15 = 0.187$  L of  $\text{H}_2$  at STP

At STP

$$P_1 = 1 \text{ atm}$$

$$V_1 = 0.187 \text{ L}$$

$$T_1 = 273 \text{ K}$$


Given conditions

$$P_2 = 1 \text{ bar} = 0.987 \text{ atm}$$

$$V_2 = ?$$

$$T_2 = 273 + 20 = 293 \text{ K}$$

# Last-minute check on your NEET readiness



₹ 500

## HIGHLIGHTS:

- 10 Model Test Papers based on latest NEET syllabus
- Last 14 years' solved test papers of AIPMT / NEET
- Includes solved NEET (Phase-1) 2016 paper
- Detailed solutions for self-assessment and to practice time management

MTG's NEET Explorer helps students self-assess their readiness for success in NEET. Attempting the tests put together by MTG's experienced team of editors and experts strictly on the NEET pattern and matching difficulty levels, students can easily measure their preparedness for success.  
**Order now!**



Scan now with your smartphone or tablet\*

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!

Applying ideal gas equation,  $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$

$$\frac{1 \times 0.187}{273} = \frac{0.987 \times V_2}{293}$$


$$V_2 = \frac{293}{0.987} \times \frac{1 \times 0.187}{273} = 0.203 \text{ L} = 203 \text{ mL}$$

38. (d):  $\text{SF}_4$  ( $sp^3d$ , trigonal bipyramidal with one equatorial position occupied by one lone pair),  $\text{CF}_4$  ( $sp^3$ , tetrahedral, no lone pair),  $\text{XeF}_4$  ( $sp^3d^2$ , square planar, two lone pairs).

39. (b):  $(\text{SiH}_3)_3\text{N}$  is planar, while  $(\text{CH}_3)_3\text{N}$  is pyramidal.

40. (a):  $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2 + \text{HBr}$    
 1-Butene   
  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$  +  $\text{CH}_3\text{CH}_2\text{CH}(\text{Br})\text{CH}_3$    
 (III) Minor   
 A (exists in 2 enantiomers I and II) Major

(A) can have two spatial arrangements :



41. (b):  $\text{CH}_3\text{COOH} \xrightarrow{\text{NaOH(aq)}(X)} \text{CH}_3\text{COONa} \xrightarrow[\Delta]{\text{sodalime}(Y)} \text{CH}_3\text{Cl} \xleftarrow[\text{(Z)}]{\text{Cl}_2/h\nu} \text{CH}_4$

42. (c): For bcc,  $r = \frac{\sqrt{3}}{4} a = \frac{\sqrt{3}}{4} \times 5 = 2.165 \text{ \AA} = 216.5 \text{ pm}$

43. (c): In case of fcc lattice, number of atoms per unit cell,  $Z = 4$  atoms

$$\text{Therefore, } M = \frac{d N_A a^3}{Z}$$

$$= \frac{8.92 \times 6.022 \times 10^{23} \times (3.608 \times 10^{-8})^3}{4} = 63.1 \text{ g mol}^{-1}$$

Atomic mass of copper = 63.1 u

44. (d): This method is adopted to protect  $-\text{NH}_2$  group and to reduce its activating influence.

45. (b):  $V_1 = 68.6 \text{ mL}$ ,  $P_1 = 756 \text{ mm}$ ,  $T_1 = 300 \text{ K}$    
  $V_2 = ?$ ,  $P_2 = 760 \text{ mm}$ ,  $T_2 = 273 \text{ K}$


$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$\text{At NTP, vol. of } \text{N}_2, V_2 = \frac{P_1 V_1}{T_1} \cdot \frac{T_2}{P_2} = \frac{756 \times 68.6}{300} \times \frac{273}{760} = 62.09 \text{ mL}$$

Percentage of nitrogen in organic compound

$$= \frac{28}{22400} \times \frac{V_2}{w} \times 100 = \frac{28}{22400} \times \frac{62.09}{0.52} \times 100 = 14.93\%$$

46. (c):  $1^\circ \text{R} - \text{NH}_2 + \text{RCHO}$  followed by reduction.




47. (a): Both NaCl and KCl have different molecular masses.

48. (d): The stability of fluorides increases as the size of central atom increases. Thus, the correct order is  $\text{PbF}_2 > \text{SnF}_2 > \text{GeF}_2$ .

49. (b): In  $[\text{Co}(\text{CN})_6]^{3-}$ , oxidation state of Co = +3

As  $\text{CN}^-$  is a strong field ligand, it results in pairing of electrons hence,

$$\text{Co}^{3+}: 3d^6 \text{ configuration} = t_{2g}^6 e_g^0$$



50. (a): It is clear from the figure that the process has been carried out in infinite steps hence, it is an isothermal reversible expansion.

$$w = -2.303 nRT \log \frac{V_2}{V_1}$$

$$\text{But, } P_1 V_1 = P_2 V_2 \Rightarrow \frac{V_2}{V_1} = \frac{P_1}{P_2} = \frac{2}{1} = 2$$

$$\therefore w = -2.303 nRT \log \frac{P_1}{P_2}$$

$$\begin{aligned} &= -2.303 \times 1 \text{ mol} \times 8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 298 \text{ K} \times \log 2 \\ &= -2.303 \times 8.314 \times 298 \times 0.3010 \text{ J} \\ &= -1717.63 \text{ J} \end{aligned}$$






# EXAM PREP


## Questions for Practice

- A flask contains  $H_2$  gas and a few drops of water. Pressure in the flask is 755 mm Hg at certain temperature  $T$  K. If the temperature is increased by 1%, the pressure inside the flask will be (Given, aqueous tension at two temperatures are 25 mm Hg and 30 mm Hg)  
(a) 762.1 (b) 767.3 (c) 770.4 (d) 775.6
- Consider the following reactions of alcohols :  
I.  $CH_2=CHCH_2OH \xrightarrow{A} CH_2=CHCHO$   
II.  $(CH_3)_3COH \xrightarrow{B} CH_3-C(CH_3)=CH_2$   
III.  $CH_3CH_2OH \xrightarrow{C} CH_3CO_2H$   
IV.  $CH_3CH=CHCH_3 \xrightarrow{D} CH_3CO_2H$   
A, B, C and D are respectively  
(a)  $MnO_2, Cu/\Delta, H_2CrO_4, KMnO_4/\Delta$   
(b)  $Cu/\Delta, MnO_2, H_2CrO_4, KMnO_4/\Delta$   
(c)  $MnO_2, Cu/\Delta, KMnO_4/\Delta, H_2CrO_4$   
(d)  $MnO_2, H_2CrO_4, Cu/\Delta, KMnO_4$
- Which of the following statements is correct?  
(a) Diamond is unaffected by conc. acids but graphite reacts with hot conc.  $HNO_3$  forming graphitic acid,  $C_6(COOH)_6$ .  
(b) CO is toxic because it forms a complex with haemoglobin in the blood.  
(c)  $C_3O_2$ , carbon suboxide, is a foul-smelling gas.  
(d) All of these
- In a cell that utilises the reaction,  
 $Zn_{(s)} + 2H^+_{(aq)} \longrightarrow Zn^{2+}_{(aq)} + H_{2(g)}$ , addition of  $H_2SO_4$  to cathode compartment will  
(a) lower the value of  $E_{cell}$  and shift equilibrium to the right  
(b) lower the value of  $E_{cell}$  and shift equilibrium to the left  
(c) increase the value of  $E_{cell}$  and shift equilibrium to the right  
(d) increase the value of  $E_{cell}$  and shift equilibrium to the left.
- Which is not the correct reaction?  
(a)  $\longrightarrow$  +  $NH_4Cl$   
(b)  $\longrightarrow$  +  $NH_4Cl$   
(c)  $\longrightarrow$  +  $NH_4Cl$   
(d)  $\xrightarrow{273\text{ K}}$
- Which of the following reactions should be most favoured thermodynamically?  
(a)  $Na_2O + Cl_2O_7 \longrightarrow 2NaClO_4$   
(b)  $Na_2O + SO_3 \longrightarrow Na_2SO_4$   
(c)  $3Na_2O + P_2O_5 \longrightarrow 2Na_3PO_4$   
(d)  $Na_2O + SiO_2 \longrightarrow Na_2SiO_3$
- The enthalpy changes for the following processes are listed below :  
 $Cl_{2(g)} \longrightarrow 2Cl_{(g)}; \quad 242.3 \text{ kJ mol}^{-1}$   
 $I_{2(g)} \longrightarrow 2I_{(g)}; \quad 151.0 \text{ kJ mol}^{-1}$   
 $I_{2(s)} \longrightarrow I_{2(g)}; \quad 62.76 \text{ kJ mol}^{-1}$   
 $ICl_{(g)} \longrightarrow I_{(g)} + Cl_{(g)}; \quad 211.3 \text{ kJ mol}^{-1}$   
Given that the standard states for iodine and chlorine are  $I_{2(s)}$  and  $Cl_{2(g)}$ , standard enthalpy of formation of  $ICl_{(g)}$  is  
(a)  $-16.8 \text{ kJ mol}^{-1}$  (b)  $+16.8 \text{ kJ mol}^{-1}$   
(c)  $+244.8 \text{ kJ mol}^{-1}$  (d)  $-14.6 \text{ kJ mol}^{-1}$
- In which case, chiral carbon is not generated?  
(a)  $CH_3COCH_3 + HCN \xrightarrow{H_3O^+}$   
(b)  $CH_3CHO + HCN \xrightarrow{H_3O^+}$   
(c)  $CH_3COCOOH + HCN \xrightarrow{H_3O^+}$   
(d)  $\xrightarrow{H_3O^+}$
- What may be expected to happen when phosphine gas is mixed with chlorine gas?  
(a)  $PH_3 \cdot Cl_2$  is formed with warming up.  
(b) The mixture only cools down.



- (c)  $\text{PCl}_3$  and  $\text{HCl}$  are formed and the mixture warms up.  
 (d)  $\text{PCl}_5$  and  $\text{HCl}$  are formed and the mixture cools down.
10. The hydrolysis of ethyl acetate was carried out separately with 0.05 M  $\text{HCl}$  and 0.05 M  $\text{H}_2\text{SO}_4$ . The rate constants were found to be  $k_1$  and  $k_2$  respectively then  
 (a)  $k_1 < k_2$  (b)  $k_1 > k_2$   
 (c)  $k_1 = k_2$  (d)  $k_2 = 2k_1$
11. The nodal plane in the  $\pi$ -bond of ethene is located in  
 (a) the molecular plane  
 (b) a plane parallel to the molecular plane  
 (c) a plane perpendicular to the molecular plane which bisects the carbon-carbon sigma bond at right angle  
 (d) a plane perpendicular to the molecular plane which contains the carbon-carbon sigma bond.
12. The water pollutants mainly responsible for eutrophication are  
 (a)  $\text{Cd}$ ,  $\text{Pb}$  and  $\text{Hg}$  present in industrial waste  
 (b) heavy metals present in mining waste  
 (c) detergents and fertilizers containing phosphate anion  
 (d) polychlorinated biphenyls.
13. If  $\lambda_0$  is the threshold wavelength of a metal and  $\lambda$  is the wavelength of the incident radiation the maximum velocity of the ejected electrons from the metal would be  
 (a)  $\left[ \frac{hc}{m} \left( \frac{\lambda - \lambda_0}{\lambda \lambda_0} \right) \right]^{1/2}$  (b)  $\left[ \frac{2hc}{m} \left( \frac{\lambda_0 - \lambda}{\lambda_0 \lambda} \right) \right]^{1/2}$   
 (c)  $\left[ \frac{h}{m} (\lambda_0 - \lambda) \right]^{1/2}$  (d)  $\left[ \frac{2h}{m} (\lambda - \lambda_0) \right]^{1/2}$
14. A new carbon-carbon bond formation is possible in  
 (a) Cannizzaro reaction  
 (b) Reimer-Tiemann reaction  
 (c) Clemmensen reduction  
 (d) none of the above.
15. The electronic spectrum of  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$  gives a single broad peak with a maxima at  $20300 \text{ cm}^{-1}$ . The crystal field stabilization energy is equal to ( $1 \text{ kJ mol}^{-1} \approx 83.7 \text{ cm}^{-1}$ )  
 (a)  $97 \text{ kJ mol}^{-1}$  (b)  $243 \text{ kJ mol}^{-1}$   
 (c)  $297 \text{ kJ mol}^{-1}$  (d) none of these.
16. Which of the following is correct, if for the same element three separate radii 105 nm, 100 nm and 90 nm are reported?  
 (a) These are van der Waals', metallic and covalent radii.  
 (b) These are metallic, covalent and van der Waals' radii respectively.  
 (c) These are covalent, metallic and van der Waals' radii respectively.  
 (d) These are metallic, van der Waals' and covalent radii respectively.
17. The compound which is not isomeric to cyclobutanol is  
 (a)  (b) butanone  
 (c) but-3-ene-1-ol (d) diethyl ether.
18.  $\text{Pb}_3\text{O}_4$  reacts with  $\text{HNO}_3$  forming nitrate and oxide which are  
 (a)  $\text{Pb}(\text{NO}_3)_2$  from  $\text{PbO}$ , and  $\text{PbO}_2$  remains unreacted  
 (b)  $\text{Pb}(\text{NO}_3)_2$  from  $\text{PbO}_2$  and  $\text{PbO}$  remains unreacted  
 (c)  $\text{Pb}(\text{NO}_3)_2$  from  $\text{Pb}(\text{NO}_3)_4$  and  $\text{PbO}$  remains unreacted  
 (d)  $\text{Pb}_3\text{O}_4$  remains insoluble in  $\text{HNO}_3$ .
19. In a particular solution of sulphur dissolved in carbon disulphide, 80% of the sulphur atoms exist as  $\text{S}_8$  while remaining as  $\text{S}_2$ . What should be the value of van't Hoff factor ' $i$ ' for the calculation of colligative properties?  
 (a) 1 (b)  $\frac{1}{2}$   
 (c)  $\frac{1}{5}$  (d)  $\frac{1}{8}$
20. How many chirality centers are present in 1-chloro-2-methylcyclohexane?  
 (a) 2 (b) 4 (c) 1 (d) 3
21. Cerium ( $Z = 58$ ) is an important member of the lanthanides. Which of the following statements about cerium is incorrect?  
 (a) The common oxidation states of cerium are +3 and +4.  
 (b) Cerium (IV) acts as an oxidising agent.  
 (c) The +4 oxidation state of cerium is not known in solutions.  
 (d) The +3 oxidation state of cerium is more stable than the +4 oxidation state.


22. ZSM-5 is one of the zeolite with formula  
 (a)  $H_x[(AlO_2)_x \cdot (SiO_2)_{96-x}] \cdot 16H_2O$   
 (b)  $Na_{56}[(AlO_2)_{56} \cdot (SiO_2)_{36}] \cdot 250H_2O$   
 (c)  $Na_2Ca(AlO_2)_2(SiO_2)_4 \cdot 6H_2O$   
 (d) none of the above is correct.
23. Which of the following hydrocarbons has the lowest dipole moment?  
 (a)  $\begin{array}{c} H_3C \\ \diagdown \\ C \\ \diagup \\ H \end{array} = \begin{array}{c} CH_3 \\ \diagup \\ C \\ \diagdown \\ H \end{array}$  (b)  $CH_3C \equiv CCH_3$   
 (c)  $CH_3CH_2C \equiv CH$  (d)  $CH_2=CH-C \equiv CH$
24. Which of the following are peroxy acids?  
 (a)  $H_2S_2O_3$  and  $H_2S_2O_6$  (b)  $H_2S_2O_6$  and  $H_2SO_5$   
 (c)  $H_2SO_5$  and  $H_2S_2O_8$  (d)  $H_2S_2O_3$  and  $H_2S_2O_8$
25. Assuming complete dissociation which of the following aqueous solutions will have the same pH value?  
 (i) 100 mL of 0.01 M HCl  
 (ii) 100 mL of 0.01 M  $H_2SO_4$   
 (iii) 50 mL of 0.01 M HCl  
 (iv) Mixture of 50 mL of 0.02 M  $H_2SO_4$  and 50 mL of 0.02 M NaOH  
 (a) (i), (ii) (b) (i), (iii)  
 (c) (ii), (iv) (d) (i), (iv)
26. Which of the following will require maximum volume of x M  $H_2SO_4$  for complete reaction?  
 (a) 65 mg of Zn  
 (b) 20 mL of 0.15 M KOH  
 (c) 15 mL of 0.2 M  $Na_2CO_3$   
 (d) 25 mL of 0.1 M  $Ba(OH)_2$
27. The acidic character of  $H_2O$ ,  $H_2O_2$  and  $CO_2$  in increasing order is  
 (a)  $H_2O < H_2O_2 < CO_2$  (b)  $H_2O < CO_2 < H_2O_2$   
 (c)  $H_2O_2 < H_2O < CO_2$  (d)  $CO_2 < H_2O < H_2O_2$
28. Molecular weight of  $H_3PO_4$  is 98. Its equivalent weight in the reaction given below is  
 $H_3PO_4 + NaOH \longrightarrow H_2O + NaH_2PO_4$   
 (a) 98 (b) 49  
 (c) 294 (d) 32.67
29. Which one of the following statements about protein structure is correct?  
 (a) Proteins consisting of one polypeptide can have quaternary structure.  
 (b) The formation of a disulphide bond in a protein requires that the two participating cysteine residues be adjacent to each other in the primary sequence of the protein.
- (c) The stability of quaternary structure in proteins is mainly due to covalent bonds among the sub-units.  
 (d) The information required for the correct folding of a protein is contained in the specific sequence of amino acids along the polypeptide chain.
30.  $BeF_2$  is soluble in water whereas fluorides of other alkaline earth metals are insoluble because of  
 (a) ionic nature of  $BeF_2$   
 (b) covalent nature of  $BeF_2$   
 (c) greater hydration energy of  $Be^{2+}$  ion as compared to its lattice energy  
 (d) none of these.
31. Consider the following equilibria at 300 K and 400 K with their equilibrium constants :
- |  | 300 K         | 400 K        |
|--|---------------|--------------|
| I. $A_{(g)} \rightleftharpoons 2B_{(g)}$ | $K_{eq} = 10$ | $K_{eq} = 5$ |
| II. $C_{(g)} \rightleftharpoons D_{(g)}$ | $K_{eq} = 2$  | $K_{eq} = 5$ |
- The data shows that  
 (a) I is endothermic, II is exothermic  
 (b) I is exothermic, II is endothermic  
 (c) I and II both are endothermic  
 (d) I and II both are exothermic.
32. Which of the following detergents acts as a germicide?  
 (a) Cetyltrimethylammonium chloride  
 (b) Sodium laurylsulphate  
 (c) Sodium-4-(1-dodecyl)benzenesulphonate  
 (d) All of the above
33. Select the incorrect statement.  
 (a)  $H^+$  can exist as  $H_9O_4^+$  in water.  
 (b)  $H_2$  is thermally stable.  
 (c) Ionisation of  $CH_3COOH$  is slower than that of  $CH_3COOD$ .  
 (d) Kinetic isotopic effect is observed when there is retardation in the rate of a reaction when  $H_2O$  is replaced by  $D_2O$ .
34. The pycnometric density of sodium chloride crystal is  $2.165 \times 10^3 \text{ kg m}^{-3}$ , while its X-rays density is  $2.178 \times 10^3 \text{ kg m}^{-3}$ . The fraction of unoccupied sites in sodium chloride crystal is  
 (a) 5.96 (b)  $5.96 \times 10^{-1}$   
 (c)  $5.96 \times 10^{-2}$  (d)  $5.96 \times 10^{-3}$
35. The major organic products A and B in the given reactions are respectively



36. Which of the following is not correct?
- Chromyl bromide test is conducted for  $\text{Br}^-$  ion.
  - Borax bead test is for coloured salts.
  - Dimethylglyoxime is a test reagent for  $\text{Ni}^{2+}$ .
  - Ring test is performed for  $\text{NO}_3^-$ .

37. Which among the following statements is false?
- Adsorption may be monolayered or multilayered.
  - Particle size of adsorbent will not affect the amount of adsorption.
  - Increase of pressure increases the amount of adsorption.
  - Increase of temperature may decrease the amount of adsorption.

38. Which of the following does not exhibit tautomerism?





39. Which one of the following sequences represents the increasing order of the polarizing power of the cationic species,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Be}^{2+}$ ?

- $\text{Mg}^{2+}$ ,  $\text{Be}^{2+}$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$
- $\text{Be}^{2+}$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$
- $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Be}^{2+}$
- $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Be}^{2+}$ ,  $\text{K}^+$

40. 1 g  $\text{H}_2$  gas at STP is expanded so that volume is doubled. Hence, work done is

- 22.4 L atm
- 5.6 L atm
- 11.2 L atm
- 44.8 L atm



42.  $\text{HNO}_3$  is changing to  $\text{N}_2\text{O}$ ,  $\text{NO}$ ,  $\text{NO}_2$  and  $\text{N}_2\text{O}_5$ . The changes involved in number of electrons per N-atom are respectively

- 0, 1, 2, 4
- 4, 3, 1, 0
- 1, 2, 4, 5
- 3, 1, 4, 0

43.  $\text{NH}_3$  gas is liquefied more easily than  $\text{N}_2$ . Hence,

- values of van der Waals' constants  $a$  and  $b$  of  $\text{NH}_3$  are greater than that of  $\text{N}_2$
- values of van der Waals' constant  $a$  and  $b$  of  $\text{NH}_3$  are less than that of  $\text{N}_2$
- $a(\text{NH}_3) > a(\text{N}_2)$  but  $b(\text{NH}_3) < b(\text{N}_2)$
- $a(\text{NH}_3) < a(\text{N}_2)$  but  $b(\text{NH}_3) > b(\text{N}_2)$



44. Which of the following statements is correct w.r.t. the reactions of aqueous solution of methanamine?

- It converts aqueous  $\text{CuSO}_4$  to blue
- It dissolves  $\text{AgCl}$  in it
- $\text{Fe}^{3+}$  gives brown precipitate of  $\text{Fe}(\text{OH})_3$
- All of the above

45. The correct order of crystal field stabilisation energies for the given ions is

- $\text{V}^{2+} < \text{Mn}^{2+} < \text{Fe}^{2+} < \text{Co}^{2+} < \text{Ni}^{2+}$
- $\text{Ni}^{2+} < \text{Co}^{2+} < \text{Fe}^{2+} < \text{Mn}^{2+} < \text{V}^{2+}$
- $\text{Mn}^{2+} < \text{V}^{2+} < \text{Co}^{2+} < \text{Fe}^{2+} < \text{Ni}^{2+}$
- $\text{V}^{2+} \approx \text{Mn}^{2+} \approx \text{Fe}^{2+} \approx \text{Co}^{2+} \approx \text{Ni}^{2+}$

46. Butanone can be best obtained by the ozonolysis of



B, C and D respectively are

- aspirin, salol, oil of wintergreen
- salol, aspirin, oil of wintergreen
- oil of wintergreen, aspirin, salol
- oil of wintergreen, salol, aspirin.


48. The colourless species is  
 (a)  $\text{VCl}_3$  (b)  $\text{VO}\text{SO}_4$   
 (c)  $\text{Na}_3\text{VO}_4$  (d)  $[\text{V}(\text{H}_2\text{O})_6]\text{SO}_4 \cdot \text{H}_2\text{O}$
49. In the reaction,  
 $2\text{Al}_{(s)} + 6\text{HCl}_{(aq)} \longrightarrow 2\text{Al}_{(aq)}^{3+} + 6\text{Cl}_{(aq)}^- + 3\text{H}_{2(g)}$   
 (a) 6 L  $\text{HCl}_{(aq)}$  is consumed for every 3 L  $\text{H}_{2(g)}$  produced  
 (b) 33.6 L  $\text{H}_{2(g)}$  is produced regardless of temperature and pressure for every mole of Al reacted  
 (c) 67.2 L  $\text{H}_{2(g)}$  at STP is produced for every mole of Al reacted  
 (d) 11.2 L  $\text{H}_{2(g)}$  at STP is produced for every mole of  $\text{HCl}_{(aq)}$  consumed.
50. Assume that a particular amino acid has an isoelectric point of 6.0. In a solution of pH 1.0, which of the following species will predominate?
- (a)  $\text{H}_3\text{N}^+\text{CH}(\text{R})\text{CO}_2\text{H}$  (b)  $\text{H}_2\text{NCH}(\text{R})\text{CO}_2\text{H}$   
 (c)  $\text{H}_3\text{N}^+\text{CH}(\text{R})\text{CO}_2^-$  (d)  $\text{H}_2\text{NCH}(\text{R})\text{CO}_2^-$

### SOLUTIONS

1. (b): Pressure of dry  $\text{H}_2$  at  $T \text{ K} = 755 - 25 = 730 \text{ mm Hg}$   
 New temperature  $= T + \frac{1}{100}T = 1.01 T$   
 New pressure of dry  $\text{H}_2 = \frac{P_1}{T_1} \times T_2 = \frac{730}{T} \times 1.01 T$   
 $= 737.3 \text{ mm Hg}$   
 New pressure of moist  $\text{H}_2$  in flask  $= 737.3 + 30$   
 $= 767.3 \text{ mm Hg}$
2. (a):  $\text{MnO}_2$  is very selective and can oxidise only allylic, benzylic or propargylic alcohols.
3. (d)
4. (c):  $\text{Zn}_{(s)} + 2\text{H}^+_{(aq)} \longrightarrow \text{Zn}^{2+}_{(aq)} + \text{H}_{2(g)}$   

$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.059}{2} \log \frac{[\text{Zn}^{2+}]}{[\text{H}^+]^2}$$
  
 When  $\text{H}_2\text{SO}_4$  is added to cathode compartment then,  $[\text{H}^+]$  will increase therefore,  $E_{\text{cell}}$  will also increase and equilibrium will shift towards right.
5. (b):  $\text{NH}_3$  (a strong nucleophile) causes elimination from  $3^\circ$  alkyl halide.
6. (a):  $\text{Na}_2\text{O}$  is basic in nature and  $\text{Cl}_2\text{O}_7$  is most acidic among the given oxides.

Greater the acidic nature, greater is the tendency of the reaction to occur thus, reaction of  $\text{Na}_2\text{O}$  with  $\text{Cl}_2\text{O}_7$  is most favourable.



$$\begin{aligned} \Delta H &= \left( \frac{1}{2} \Delta_{\text{sub}} H + \frac{1}{2} \Delta_{\text{diss}} H \right)_{\text{I}_2} + \frac{1}{2} (\Delta_{\text{diss}} H)_{\text{Cl}_2} + \Delta H' \\ &= \frac{1}{2} \times 151 + \frac{1}{2} \times 62.76 + \frac{1}{2} \times 242.3 - 211.3 \\ &= 75.5 + 31.38 + 121.15 - 211.3 \\ &= 228.03 - 211.3 = 16.73 \text{ kJ mol}^{-1} \end{aligned}$$

8. (a)
9. (d):  $\text{PH}_3 + 4\text{Cl}_2 \longrightarrow \text{PCl}_5 + 3\text{HCl}$   
 $\text{Cl}_2$  oxidises  $\text{P}^{3-}$  to  $\text{P}^{5+}$ . The reaction is endothermic thus, the mixture cools down.
10. (a): In the presence of an acid, hydrolysis of ethyl acetate is a pseudo-unimolecular reaction but the actual value of  $k$  depends upon the concentration of  $\text{H}^+$  ions. As  $\text{H}_2\text{SO}_4$  is a stronger acid than  $\text{HCl}$  and moreover  $\text{H}^+$  ions produced from 0.05 M  $\text{H}_2\text{SO}_4$  is double than 0.05 M  $\text{HCl}$  therefore,  $k_2 > k_1$ .
11. (a)      12. (c)
13. (b):  $E = K.E. + E_{\text{threshold}}$   

$$\frac{hc}{\lambda} = \frac{1}{2}mv^2 + \frac{hc}{\lambda_0}$$
  

$$\frac{1}{2}mv^2 = \frac{hc}{\lambda} - \frac{hc}{\lambda_0}$$
  

$$v^2 = \frac{2hc}{m} \left( \frac{1}{\lambda} - \frac{1}{\lambda_0} \right)$$
  

$$v = \left[ \frac{2hc}{m} \left( \frac{\lambda_0 - \lambda}{\lambda_0 \lambda} \right) \right]^{1/2}$$
14. (b): In the Reimer-Tiemann reaction, new C—C bond is formed between the carbon of benzene ring and —CHO group.

# CONCEPT MAP

## ALDEHYDES, KETONES AND CARBOXYLIC ACIDS

Being important constituents of fabrics, flavourings, plastics and drugs, carbonyl compounds are of utmost importance to organic chemistry while carboxylic acids are amongst the earliest organic compounds to be isolated from nature and are still known by their common names.

### Structure and Nomenclature

#### Aldehydes

- $R-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-\text{H}$  where,  $R = \text{H}$ , alkyl or aryl group.
- In IUPAC system, aldehydes are named as *alkanals*.

#### Ketones

- $R-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-R'$  where  $R$  and  $R'$  both can be same or different groups.
- In IUPAC system, they are named as *alkanones*.

### ALDEHYDES AND KETONES

### Physical Properties

- Solubility in water  $\propto \frac{1}{\text{Molecular mass}}$
- Compounds having upto four carbon atoms are soluble in water due to hydrogen bonding.
- Due to dipole-dipole interactions their b.p.s. are higher than the corresponding hydrocarbons or ethers but lesser than alcohols or carboxylic acids which have intermolecular H-bonding.
- Due to two electron donating alkyl groups, ketones have higher b.p.s. than the corresponding aldehydes.

### Distinction Tests

Test	Aldehydes	Ketones
Schiff's reagent	Pink colour	No colour
Fehling's solution	Red ppt.	No ppt.
Tollens' reagent	Silver mirror	No ppt.
Sodium hydroxide	Brown resinous mass (except HCHO)	No reaction
Alkaline sodium nitroprusside	A deep red colour (except HCHO)	Red colour which changes to orange

### Structure and Nomenclature

#### Carboxylic acids

- $R-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-\text{OH}$  where,  $R = \text{H}$ , alkyl or aryl group.
- In IUPAC system, they are named as *alkanoic acids*.

### Physical Properties

- Solubility in water  $\propto \frac{1}{\text{Molecular mass}}$
- High b.pt. due to intermolecular hydrogen bonding.
- M.p.s. and b.p.s. of aromatic acids are usually higher than those of aliphatic acids.

### CARBOXYLIC ACIDS

### Chemical Properties

- Acidity order:** Carboxylic acids > Phenols > Alcohols
- EDG decreases the acidity and EWG increases the acidity.
- More the electronegativity of the atom attached to the carboxyl group, more will be the acidity.

### Distinction Tests

Test	Carboxylic acids	Phenols	Alcohols
$\text{NaHCO}_3$	Brisk effervescence of $\text{CO}_2$ gas	No reaction	No reaction
$\text{FeCl}_3$	Buff coloured ppt.	Violet, blue or red colour	No reaction

### Preparation

- Oxidation of alcohols:**

$$\text{RCH}_2\text{OH} + [\text{O}] \xrightarrow{\text{K}_2\text{Cr}_2\text{O}_7/\text{H}_2\text{SO}_4(\text{dil.})} \text{RCHO} + \text{H}_2\text{O}$$

(1° alcohol)

$$\text{RCH}(\text{OH})\text{R}' + [\text{O}] \xrightarrow{\text{K}_2\text{Cr}_2\text{O}_7/\text{H}_2\text{SO}_4(\text{dil.})} \text{RCOR}' + \text{H}_2\text{O}$$


(2° alcohol)
- Catalytic decomposition of carboxylic acids:**

$$\text{RCOOH} + \text{HOOCH} \xrightarrow{\text{MnO}, 573 \text{ K}} \text{RCHO} + \text{CO}_2 + \text{H}_2\text{O}$$

$$\text{RCOOH} + \text{HOOCR}' \xrightarrow{\text{MnO}, 573 \text{ K}} \text{RCOR}' + \text{CO}_2 + \text{H}_2\text{O}$$
- Hydroboration-oxidation of alkynes:**

$$\text{R}-\text{C}\equiv\text{C}-\text{H} \xrightarrow[\text{THF}]{\text{B}_2\text{H}_6} \left[ \begin{array}{c} \text{R}-\text{C}=\text{C}-\text{H} \\ | \quad | \\ \text{H} \quad \text{BH}_2 \end{array} \right] \xrightarrow[\text{OH}^-]{\text{H}_2\text{O}_2} \text{R}-\text{C}(\text{OH})=\text{C}-\text{H} \rightleftharpoons \text{RCH}_2\text{C}(=\text{O})-\text{H}$$


Terminal alkyne



- Ozonolysis of alkenes:**


$$\text{RCH}=\text{C}(\text{R}')-\text{R}'' \xrightarrow[\text{H}_2\text{O}, \text{Zn}]{\text{O}_3} \text{RCHO} + \text{R}'\text{C}(=\text{O})\text{R}''$$

- From gem-dihalides:**



### Chemical Properties


- Nucleophilic addition reactions:**





- Reactivity order:**

Aldehydes > Ketones (steric and electronic reasons)  
 $\text{HCHO} > \text{RCHO} > \text{PhCHO} > \text{RCOR} > \text{RCOPh} > \text{PhCOPh}$


- Nucleophilic addition-elimination reactions:**




- Reduction and oxidation:**




- Haloform reaction:**



- Aldol condensation:**



- Cannizzaro reaction:**



### Preparation

- Oxidation of 1° alcohols:**


$$\text{RCH}_2\text{OH} \xrightarrow[\text{[O]}]{\text{alk. KMnO}_4} \text{RCHO} \xrightarrow[\text{acid}]{\text{[O]}} \text{RCOOH}$$
- Hydrolysis of nitriles and amides:**

$$\text{RCN} + 2\text{H}_2\text{O} \xrightarrow[\text{OH}^-]{\text{H}^+ \text{ or } \text{OH}^-} \text{RCOOH} + \text{NH}_3$$


$$\text{RCONH}_2 \xrightarrow[\text{heat}]{\text{H}_2\text{O}} \text{RCOOH} + \text{NH}_3$$
- From Grignard reagents:**

$$\text{CO}_2 + \text{CH}_3\text{MgBr} \xrightarrow[\text{H}^+, \text{H}_2\text{O}]{\text{Dry ether}} \text{CH}_3\text{COOH} + \text{Mg(OH)Br}$$


- Hydrolysis of esters:**



- Carbonylation (Koch reaction) of alkenes:**




- From methyl ketones:**




### Chemical Reactions


- Reactions involving cleavage of  $-\text{OH}$  group:**



- Reactions involving proton of  $-\text{OH}$  group:**




- Reactions involving  $>\text{C}=\text{O}$  group:**




- Ring substitution in aromatic acids:**


$-\text{COOH}$  group is deactivating and *meta* directing.




- Reactions involving  $-\text{COOH}$  group:**



- Reactions involving  $-\text{R}$  group:**








15. (b):  $1 \text{ kJ mol}^{-1} = 83.7 \text{ cm}^{-1}$

$$\Delta_o = \frac{20300}{83.7} = 243 \text{ kJ mol}^{-1}$$


16. (a): van der Waals' radius > metallic radius > covalent radius.

17. (d): Cyclobutanol is  ( $\text{C}_4\text{H}_8\text{O}$ ), diethyl ether is  $\text{C}_2\text{H}_5\text{OC}_2\text{H}_5$  ( $\text{C}_4\text{H}_{10}\text{O}$ ).

18. (a):  $\text{Pb}_3\text{O}_4 + 4\text{HNO}_3 \longrightarrow 2\text{Pb}(\text{NO}_3)_2 + \text{PbO}_2 + 2\text{H}_2\text{O}$

19. (c):  $8 \text{ S} \longrightarrow \text{S}_8$

0.8 mol of S gives 0.1 mol of  $\text{S}_8$




0.2 mol of S gives 0.1 mol of  $\text{S}_2$

Total moles =  $0.1 + 0.1 = 0.2$

$\therefore$  1 mole of sulphur atoms behave as 0.2 moles.

So,  $i = 0.2$


20. (a): A carbon bonded with four different types of groups or atoms is known as chiral carbon.



21. (c): +4 oxidation state of cerium is also known in solutions.

22. (a)

23. (b):  $\text{CH}_3-\text{C}\equiv\text{C}-\text{CH}_3$ . It is because symmetrical individual dipoles get cancelled leading to least dipole moment. Others are unsymmetrical, therefore, dipoles do not get cancelled.



25. (d): Number of eq. of  $0.01 \text{ M HCl} = \frac{0.01 \times 100}{1000} = 1 \times 10^{-3}$

$$\text{pH} = -\log[1 \times 10^{-3}] = 3$$

$$\begin{array}{l} \text{Number of eq. of } 0.02 \text{ M H}_2\text{SO}_4 = \frac{0.04 \times 50}{1000} \\ = 2 \times 10^{-3} \end{array}$$

$$\begin{array}{l} \text{Number of eq. of } 0.02 \text{ M NaOH} = \frac{0.02 \times 50}{1000} \\ = 1 \times 10^{-3} \end{array}$$

$$[\text{H}^+] \text{ left} = 2 \times 10^{-3} - 1 \times 10^{-3} = 1 \times 10^{-3} : \text{pH} = 3$$

26. (c): Millimoles of  $\text{H}_2\text{SO}_4$  required for different reagents are as follows:

$$\text{For Zn} = \frac{65}{65} = 1$$

$$\text{For KOH} = \frac{20 \times 0.15}{2} = 1.5$$

$$\text{For Na}_2\text{CO}_3 = 15 \times 0.2 = 3$$

$$\text{For Ba(OH)}_2 = 25 \times 0.1 = 2.5$$

27. (a)

28. (a): Only one H of  $\text{H}_3\text{PO}_4$  has been replaced. Hence,

$$\begin{array}{l} \text{Eq. wt. of H}_3\text{PO}_4 = \frac{\text{Mol wt.}}{\text{Basicity in the reaction}} \\ = \frac{98}{1} = 98 \end{array}$$

29. (d): Quaternary structure is observed in proteins which exists as assemblies of two or more polypeptide chains called sub-units which are held together by non-covalent forces such as hydrogen bonds, electrostatic interactions and van der Waals' interactions. The correct folding of a protein is guided by specific interactions among the side chains of the amino acid residues of a polypeptide chain. The two cysteine residues that react to form the disulphide bond may be a great distance apart in the primary structure (or on separate polypeptides) but are brought into close proximity by the three dimensional folding of the polypeptide chain.

30. (c)

31. (b): For exothermic reactions on increasing temperature, equilibrium constant gets decreased and inverse is true for endothermic reactions.


32. (a): Quaternary ammonium halides made from long chain aliphatic amines are called invert soaps because their positive ion acts as surfactant. These are used as germicides.

33. (c): Deuterium bond is stronger than the proton bond hence, ionisation of  $\text{CH}_3\text{COOD}$  is slower than  $\text{CH}_3\text{COOH}$ .

34. (d): Difference =  $2.178 \times 10^3 - 2.165 \times 10^3 = 0.013 \times 10^3$


$$\text{Fraction unoccupied} = \frac{0.013 \times 10^3}{2.178 \times 10^3} = 5.96 \times 10^{-3}$$

**35. (a) :** Dehydrohalogenation using  $\text{KOH}/\text{C}_2\text{H}_5\text{OH}$  produces Saytzeff product as the major product while the strong bases like  $\text{Et}_3\text{N}$  or  $(\text{CH}_3)_3\text{COH}/\text{KOH}$  produce Hofmann product as the major product.



**36. (a):** There is no test named as chromyl bromide test. Chromyl bromide is not stable at all.



**37. (b):** Adsorption increases with increase in surface area.



**39. (c):** Smaller the size of cation, higher will be its polarizing power.

**40. (c):** 1 g H<sub>2</sub> at STP = 0.5 mol = 11.2 L = V<sub>1</sub>  
 $W = p(V_2 - V_1) = p(2V_1 - V_1)$   
 $= pV_1 = 1 \times 11.2 \text{ L atm}$

**41. (b):**





43. (a): If a gas is liquefied, it means volume is decreased and attractive forces exist.  $a$  is a measure of attractive forces and  $b$  is the measure of excluded volume hence,

$$a(\text{NH}_3) > a(\text{N}_2) \text{ and } b(\text{NH}_3) > b(\text{N}_2)$$

44. (c)

**45. (b):**  $\text{CFSE} \propto \frac{1}{\text{Number of } d\text{-electrons}}$   
(For metal ions having same charge)



**48. (c) :** In  $\text{Na}_3\text{VO}_4$ , oxidation state of V is +5, i.e.,  $3d^0 4s^0$  electronic configuration. Hence, no  $d-d$  transition takes place due to which it is colourless.

**49. (d):**  $2\text{Al}_{(s)} + 6\text{HCl}_{(aq)} \longrightarrow 2\text{Al}^{3+}_{(aq)} + 6\text{Cl}^{-}_{(aq)} + 3\text{H}_{2(g)}$

6 mol3 × 22.4 L


3 × 22.4 L H<sub>2(g)</sub> at STP are produced from 6 moles  
HCl. Hence, 11.2 L H<sub>2(g)</sub> at STP are produced from  
1 mole HCl.

**50. (a) :** In acidic medium  $-\text{COO}^-$  group acts as a base and accepts a proton.


# OLYMPIAD PROBLEMS



1. Which molecule will give following dicarboxylic acid upon treatment with acidic solution of  $\text{KMnO}_4$ ?



2. At what temperature will hydrogen molecules have the same kinetic energy as nitrogen molecules have at  $35^\circ\text{C}$ ?



3. Which of the following pairs of molecules have the identical bond dissociation energies?



4. If the dispersed phase in colloidal iron (III) hydroxide and colloidal gold is positively and negatively charged respectively then which of the following statements is not correct?

- (a) Magnesium chloride solution coagulates gold sol readily than iron (III) hydroxide sol.
- (b) Sodium sulphate solution causes coagulation in both sol.

- (c) Mixing of the two sols has no effect.
- (d) Coagulation in both sols can be brought about by electrophoresis.


5. Consider a  $p_y$ -orbital of an atom and identify the correct statement.

- (a)  $s$ -orbital of another atom produces  $\pi$ -bond when  $y$  is the bond formation axis.
- (b)  $p_y$ -orbital of another atom produces  $\sigma$ -bond when  $x$  is the bond formation axis.
- (c)  $p_z$ -orbital of another atom produces  $\pi$ -bond when  $x$  is the bond formation axis.
- (d)  $d_{xy}$ -orbital of another atom produces  $\pi$ -bond when  $x$  is the bond formation axis.

6. Calculate the % of free  $\text{SO}_3$  in an oleum (considered as a solution of  $\text{SO}_3$  in  $\text{H}_2\text{SO}_4$ ) that is labelled '109%  $\text{H}_2\text{SO}_4$ '.

- (a) 30% (b) 40%
- (c) 35% (d) 20%

7. Among the following, which intermediate is the best hydride donor in a Cannizzaro reaction?




8. A pale green crystalline metal salt of ' $M$ ' dissolves freely in water. On standing, it gives a brown ppt. on addition of aqueous  $\text{NaOH}$ . The metal salt solution also gives a black ppt. on bubbling  $\text{H}_2\text{S}$  in basic medium. An aqueous solution of the metal salt decolourises the pink colour of the permanganate solution. The metal ' $M$ ' in the metal salt solution is

- (a) copper (b) aluminium  
(c) lead (d) iron.

9. For the equilibrium  $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$ ,  
 $K_p = \frac{\alpha^2}{(1-\alpha)V}$ ; temperature remaining constant,

- (a)  $K_p$  will increase with the increase in volume  
(b)  $K_p$  will increase with the decrease in volume  
(c)  $K_p$  will not change with the change in volume  
(d)  $K_p$  may increase or decrease with the change in volume.


10. The product of the following reaction is



- (a)
- (b)
- (c)

- (d) none of these.

11. Acetone exhibits keto-enol tautomerism as shown below:



Which of the following products is obtained when acetone is treated with an excess of  $\text{D}_2\text{O}$  for a sufficient time in the presence of a small amount of a dilute  $\text{NaOH}$  solution?

- (a)  $\text{CH}_3-\overset{\text{OD}}{\underset{|}{\text{C}}}=\text{CH}_2$  (b)  $\text{CH}_2\text{D}-\overset{\text{OH}}{\underset{|}{\text{C}}}=\text{CH}_2$   
(c)  $\text{CD}_3-\overset{\text{O}}{\underset{||}{\text{C}}}-\text{CD}_3$  (d)  $\text{CH}_3-\overset{\text{OD}}{\underset{|}{\text{C}}}=\text{CHD}$

12. A fuel oil contains significant quantity of sulphur. When the oil is burnt, the sulphur is oxidised to

$\text{SO}_2$  as;  $\text{S} + \text{O}_2 \rightarrow \text{SO}_2$ . In a city, 465 tonnes of  $\text{SO}_2$  are emitted by power plants each day. If 50% of  $\text{SO}_2$  comes from the combustion of fuel oil that contains 3% S by weight, how many tonnes of oil is burnt per day?

- (a) 3875 tonnes (b) 7750 tonnes  
(c) 1938 tonnes (d) 1140 tonnes

13. Which is the major product formed in the following reaction?



- (a)  $\text{C}_6\text{H}_5\text{CH}_2\text{CH}(\text{CH}_3)_2$   
(b)  $\text{C}_6\text{H}_5-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\underset{\text{CH}_3}{|}{\text{C}}}}-\text{CH}_3$   
(c)  $\text{C}_6\text{H}_5-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\underset{\text{CH}_3}{|}{\text{C}}}}-\text{C}_6\text{H}_5$

- (d) A mixture of all of these.

14. Which of the following represents the crystal field splitting energy diagram in a linear field (assuming that the ligands lie on the z-axis)?

- (a)  $\begin{matrix} \overline{d_{z^2}} & \overline{d_{xz}} & \overline{d_{yz}} \\ \overline{d_{xy}} & \overline{d_{x^2-y^2}} \end{matrix}$  (b)  $\begin{matrix} \overline{d_{xz}} & \overline{d_{yz}} \\ \overline{d_{xy}} & \overline{d_{x^2-y^2}} \end{matrix}$   
(c)  $\begin{matrix} \overline{d_{z^2}} \\ \overline{d_{xz}} & \overline{d_{yz}} \\ \overline{d_{xy}} & \overline{d_{x^2-y^2}} \end{matrix}$

- (d) No splitting in linear field is possible.

15. When pressure on 10 g of hydrogen is reduced from 20 to 1 atm at a constant temperature of 273 K. The gas behaves ideally. Calculate heat ' $q$ ' absorbed by the system.


- (a) 4140 cal (b) 8180 cal  
(c) 2140 cal (d) 5490 cal

16. The vapours of Hg absorb some electrons accelerated by a potential difference of 4.5 volts as a result of which light is emitted. If the full energy of single incident electron is supposed to be converted into light emitted by single Hg atom, find the wave number of the light.


- (a)  $7.209 \times 10^{-19} \text{ m}^{-1}$  (b)  $3.63 \times 10^6 \text{ m}^{-1}$   
(c)  $4.20 \times 10^4 \text{ m}^{-1}$  (d)  $5.65 \times 10^6 \text{ m}^{-1}$

17. Ammonia reacts with Nessler's reagent to give a brown precipitate known as iodide of Millon's base.

This base is believed to have the structure



18. The nitrogen atom in each of the following tertiary amines may be removed as trimethyl amine by repeated Hofmann elimination. Which of the following amines requires the greater number of Hofmann sequence to accomplish this?



19. Cyclopentadiene is unusually acidic for a hydrocarbon. This is because
- the carbon atoms of cyclopentadiene are  $sp$ -hybridised
  - cyclopentadiene is aromatic
  - the removal of a proton from it yields an aromatic anion
  - cyclopentadiene yields a highly stable free radical.

20. The kinetics of decomposition of  $\text{N}_2\text{O}_5$  in  $\text{CCl}_4$  solution is studied by measuring the evolved oxygen. If 24 mL of the gas was evolved in one hour while 35 mL of the gas was evolved when no more oxygen was coming out, calculate the fraction of  $\text{N}_2\text{O}_5$  decomposed in one hour.
- 0.486
  - 1.642
  - 0.686
  - 0.721

21. Which of the following does not have dangling bond?
- Graphite
  - Silica
  - Fullerene ( $\text{C}_{60}$ )
  - None of these

22. 36.4 g of 1, 1, 2, 2-tetrachloropropane was heated with zinc dust and the product was bubbled through ammoniacal  $\text{AgNO}_3$ . What is the weight of precipitate obtained?
- 25.3 g
  - 42.6 g
  - 8.0 g
  - 29.4 g


23. Gold and platinum dissolve in aqua regia to produce respectively

- $\text{H}[\text{AuCl}_4]$  and  $\text{H}[\text{PtCl}_3]$
- $\text{H}[\text{AuCl}_4]$  and  $\text{H}_2[\text{PtCl}_6]$
- $\text{H}_2[\text{AuCl}_6]$  and  $\text{H}[\text{PtCl}_4]$
- $\text{H}_2[\text{AuCl}_6]$  and  $[\text{PtCl}_6]$

24. Find the pH of a solution prepared by mixing 25 mL of a 0.5 M solution of  $\text{HCl}$ , 10 mL of a 0.5 M solution of  $\text{NaOH}$  and 15 mL of water.
- 0.623
  - 0.824
  - 0.912
  - 0.412

25. The osmotic pressure of an aqueous solution of sucrose is 2.47 atm at 303 K and the molar volume of the water present is  $18.10 \text{ cm}^3$ . Calculate the elevation in boiling point of the solution. Given  $\Delta H_{\text{vap}} = 540 \text{ cal/g}$ . (Assume volume of solvent equal to volume of solution.)
- $9.93 \times 10^{-2}$
  - $4.93 \times 10^{-2}$
  - $5.14 \times 10^{-2}$
  - $9.98 \times 10^{-2}$

26. 2, 6-Dimethylphenyl allyl ether, in which the carbon atom next to the oxygen in allyl group is  $^{14}\text{C}$ , on being heated to  $200^\circ\text{C}$ , gives



**BUY ONLINE**


Now you can buy

**MTG Books & Magazines**

**Log on to : [www.mtg.in](http://www.mtg.in)**



27. Which systematic diagram represents the correct chemical relations between calcium and its compounds?



28. Which of the following statements is incorrect in the context of silicones?

- (a) They are more stable to heat than other polymers.  
 (b) They are strongly water-repellent, good electrical insulators, and have non-sticking and antifoaming properties.


(c) The Si—O bond energy is higher than C—C bond energy.

(d) The Si—O bond energy is lower than C—C bond energy.

29. A galvanic cell is composed of two hydrogen electrodes, one of which is a standard one. In which of the following solutions should the other electrode be immersed to get maximum e.m.f?

- (a) 0.1 M HCl (b) 0.1 M CH<sub>3</sub>COOH  
 (c) 0.1 M H<sub>3</sub>PO<sub>4</sub> (d) 0.1 M H<sub>2</sub>SO<sub>4</sub>

30. The correct decreasing order of relative reactivity of the following chlorides toward aqueous KOH solution is




- (a)  $P > Q > R > S$  (b)  $R > P > Q > S$   
 (c)  $S > R > Q > P$  (d)  $R > S > Q > P$

## ANSWER KEYS

1. (d) 2. (d) 3. (c) 4. (c) 5. (d)  
 6. (b) 7. (d) 8. (d) 9. (c) 10. (b)  
 11. (c) 12. (a) 13. (b) 14. (c) 15. (c)  
 16. (b) 17. (a) 18. (a) 19. (c) 20. (c)  
 21. (c) 22. (d) 23. (b) 24. (b) 25. (c)  
 26. (a) 27. (a) 28. (d) 29. (b) 30. (b)

## AVAILABLE BOUND VOLUMES

of your favourite magazines



buy online at [www.mtg.in](http://www.mtg.in)

Volumes of the following years are available:

Chemistry Today

2015, 2010

Mathematics Today

2015, 2014

Biology Today

2015

Price:

₹ 325 for each volume

**How to order :** Send money by demand draft/money order. Demand Draft should be drawn in favour of MTG Learning Media (P) Ltd. Mention the volume you require along with your name and address.

**Mail your order to :**

Circulation Manager, MTG Learning Media (P) Ltd.  
 Plot 99, Sector 44 Institutional Area, Gurgaon, (HR)  
 Tel.: (0124) 6601200  
 E-mail : [info@mtg.in](mailto:info@mtg.in) Web : [www.mtg.in](http://www.mtg.in)

# ACE

YOUR WAY **CBSE XII**



Series 1

## CHAPTERWISE PRACTICE PAPER : THE SOLID STATE | SOLUTIONS

Time Allowed : 3 hours

Maximum Marks : 70

### GENERAL INSTRUCTIONS

- (i) All questions are compulsory.
- (ii) Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Q. no. 23 is a value based question and carries 4 marks.
- (vi) Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables if necessary, use of calculators is not allowed.

1. What are pseudo solids?
2. What is a de-icing agent? How does it function?
3. Give the significance of a 'lattice point'.
4. Why is  $\text{Fe}_3\text{O}_4$  ferrimagnetic at room temperature but becomes paramagnetic at 850 K?
5. Solid A is a very hard electrical insulator in solid as well as in molten state and melts at extremely high temperature. What type of solid is it?
6. In spite of long range order in the arrangement of particles, why are the crystals usually not perfect?
7. Explain with suitable examples in each case why the molar masses of some substances determined with the help of colligative properties are
  - (i) higher
  - (ii) lower than the actual values.
8. An element crystallises in a *fcc* lattice having edge length 400 pm. Calculate the maximum diameter of atom which can be placed in interstitial site without distorting the structure.
  - (i) What type of semiconductor is obtained when silicon is doped with boron?
  - (ii) What type of magnetism is shown in the following alignment of magnetic moments?  
 $\uparrow \uparrow \uparrow \uparrow \uparrow \uparrow$
9.  $\text{H}_2\text{S}$ , a toxic gas with rotten egg like smell, is used for qualitative analysis. If the solubility of  $\text{H}_2\text{S}$  in water at STP is 0.195 m, calculate Henry's law constant.
10. Schottky defects generate an equal number of cation and anion vacancies while doping produces only cation vacancies and not anion vacancies. Why?
11. (i) Niobium crystallises in body-centred cubic structure. If density is  $8.55 \text{ g cm}^{-3}$ , calculate atomic radius of niobium using its atomic mass 93 u.  
(ii) Name the parameters that characterise a unit cell.
12. Henry's law constant for  $\text{CO}_2$  in water is  $1.67 \times 10^8 \text{ Pa}$  at 298 K. Calculate the quantity of  $\text{CO}_2$  in 500 mL of soda water when packed under 2.5 atm  $\text{CO}_2$  pressure at 298 K.

OR

13. (i) Zinc oxide is white but it turns yellow on heating. Explain.  
 (ii) A sample of drinking water was found to be severely contaminated with chloroform,  $\text{CHCl}_3$  which is supposed to be a carcinogen. The level of contamination was 15 ppm (by mass).  
 (a) Express this in percent by mass.  
 (b) Determine the molality of chloroform in the water sample.
14. At  $25^\circ\text{C}$ , the vapour pressures of benzene ( $\text{C}_6\text{H}_6$ ) and toluene ( $\text{C}_7\text{H}_8$ ) are 93.4 torr and 26.9 torr, respectively. A solution is made by mixing 35.0 g of benzene and 65.0 g of toluene. At what applied pressure, in torr, will this solution boil at  $25^\circ\text{C}$ ?
15. (i) Why glasspanes fixed to windows and doors of old buildings are found to be slightly thicker at the bottom part?  
 (ii) Why the defects of crystalline solids are called thermodynamic defects?  
 (iii) What structural changes are observed when sodium chloride crystals are subjected to high pressure?
- OR**
- Metallic gold crystallises in a face-centred cubic lattice. The length of the cubic unit cell is  $a = 4.070 \text{ \AA}$ . (atomic mass of gold = 197 u)
- (i) What is the closest distance between gold atoms?  
 (ii) How many "nearest neighbours" does each gold atom have?  
 (iii) What is the density of gold?
16. (i) Osmotic pressure of a solution containing 7 g of dissolved protein per  $100 \text{ cm}^3$  of solution is 25 mm Hg at body temperature (310 K). Calculate the molecular mass of protein ( $R = 0.08205 \text{ L atm mol}^{-1} \text{ deg}^{-1}$ ).  
 (ii) Calculate the boiling point of a solution containing 0.456 g of camphor (molar mass =  $152 \text{ g mol}^{-1}$ ) dissolved in 31.4 g of acetone (b.p. =  $56.30^\circ\text{C}$ ), if the molal elevation constant of acetone is  $1.72^\circ\text{C kg mol}^{-1}$ .
17. Calculate the molarity of a solution of  $\text{CaCl}_2$ , if it is found that  $200 \text{ cm}^3$  of  $\text{CaCl}_2$  solution contains  $3.01 \times 10^{22}$  chloride ions. Assume that  $\text{CaCl}_2$  is completely ionised in solution.
18. (i) Graphite is a conductor of electricity. Give reasons.  
 (ii) How will you differentiate between *hcp* and *ccp* lattice?
19. What will be the osmotic pressure of a 0.1 M monobasic acid, if its pH is 2.0 at  $25^\circ\text{C}$ ?
20. How many mL of 0.1 M HCl are required to react completely with 1 g mixture of  $\text{Na}_2\text{CO}_3$  and  $\text{NaHCO}_3$  containing equimolar amounts of the two?
21. Aluminium crystallises in a cubic close packed structure. Radius of the atom in the metal is 125 pm.  
 (i) What is the length of the unit cell?  
 (ii) How many unit cells are there in  $1 \text{ cm}^3$  of aluminium?
22. (i) Based on solute-solvent interactions arrange the following in order of increasing solubility in *n*-octane and explain.  
 Cyclohexane, KCl,  $\text{CH}_3\text{OH}$ ,  $\text{CH}_3\text{CN}$   
 (ii) Two elements A and B form compounds having molecular formula  $\text{AB}_2$  and  $\text{AB}_4$ . When dissolved in 20 g of benzene ( $\text{C}_6\text{H}_6$ ), 1 g of  $\text{AB}_2$  lowers the freezing point by 2.3 K whereas 1 g of  $\text{AB}_4$  lowers it by 1.3 K. The molal depression constant for benzene is  $5.1 \text{ K kg mol}^{-1}$ . Calculate atomic masses of A and B.
23. Aamir, a science student dismantled an old electronic gadget to know about its components. He found that the gadget is composed of various chips, wires and other small components fitted on a circuit plate. To further explore, he searched about these chips and other components on internet and found that majority of these components are made up of a specific class of solids.  
 (i) What are the values shown by Aamir?  
 (ii) What is this class of solids called?  
 (iii) Explain why this class of solids is used to prepare components of electric devices?  
 (iv) What is the conductivity range of this class of solids?
24. (i) Heating crystals of KCl in potassium vapours makes crystals violet, why?  
 (ii) The density of KBr is  $2.75 \text{ g cm}^{-3}$ . The length of edge of the unit cell is 654 pm. Predict the type of cubic lattice to which unit cell of KBr belongs. ( $N_A = 6.023 \times 10^{23} \text{ mol}^{-1}$ ,  
 Atomic mass : K = 39, Br = 80)

- (iii) Sodium has a *bcc* structure with nearest neighbour distance 365.9 pm. Calculate its density (Atomic mass of sodium = 23).

OR

- (i) Refractive index of a solid is observed to have the same value along all directions. Comment on the nature of this solid. Would it show cleavage property?
- (ii) (a) Sodium chloride is insulator in solid state but conductor in aqueous solution. Explain why?  
(b) What kind of magnetic properties are exhibited by ions of NaCl?  
(c) What type of defects are generally observed in sodium chloride?
25. (i) Why is camphor preferred as a solvent in determination of depression in freezing point ( $\Delta T_f$ )?
- (ii) Aquatic species are more comfortable in cold water rather than in warm water. Why?
- (iii) The vapour pressures of pure liquids A and B are 450 and 700 mm Hg at 350 K respectively. Find out the composition of the liquid mixture, if total vapour pressure is 600 mm Hg. Also, find the composition of the vapour phase.

OR

- (i) When kept in water, raisins swell in size. Name and explain the phenomenon involved with the help of a diagram. Give three applications of the phenomenon.
- (ii) Explain why on addition of 1 mol of NaCl to 1 litre of water, the boiling point of water increases, while addition of 1 mol of methyl alcohol to one litre of water decreases its boiling point.
26. (i) Frenkel defects are not found in alkali metals. Explain.
- (ii)  $\text{Br}^-$  ions form close packed structure. If the radius of  $\text{Br}^-$  ion is 195 pm, calculate the radius of the cation that just fits in the tetrahedral hole. Can a cation  $\text{A}^+$  having a radius of 82 pm be slipped into the octahedral hole of the crystal  $\text{A}^+\text{Br}^-$ .
- (iii) (a) Name the non-stoichiometric point defect responsible for colour in alkali metal halides.  
(b) What is Piezoelectric effect? Explain with suitable example.

OR

- (i) In terms of band theory, what is the difference  
(a) between a conductor and an insulator  
(b) between a conductor and a semiconductor?
- (ii) Non-stoichiometric cuprous oxide,  $\text{Cu}_2\text{O}$  can be prepared in the laboratory. In this oxide, copper to oxygen ratio is slightly less than 2:1. Can you account for the fact that this substance is a *p*-type semiconductor?

## SOLUTIONS

- Amorphous solids which have tendency to flow slowly and resemble liquids in many respects are called pseudo solids or super cooled liquids. *e.g.*, glass.
- De-icing agent is a substance which prevents the formation of ice by lowering the freezing point of water to such an extent that it does not freeze to form ice.
- Each lattice point represents one constituent particle of the solid which may be an atom, an ion, or a molecule.
- $\text{Fe}_3\text{O}_4$  is ferrimagnetic at room temperature because in its crystals magnetic domains of Fe (II) and Fe (III) ions are unequal in magnitude and aligned in parallel and antiparallel directions but when heated at 850 K, the arrangement is randomised and substance becomes paramagnetic.
- Covalent or network solid.
- An ideal crystal is one which has the same unit cell containing the same lattice points across the whole crystal. But when crystallisation of solids takes place specially at fast or moderate rate some deviations from ideal arrangement may be introduced which are called defects. Thus, crystals are usually imperfect.
- (i) When a solute undergoes association in the solution then the number of solute particles decreases. Hence, the colligative property will be lower and the molar mass will be higher *e.g.*,  $2\text{CH}_3\text{COOH} \longrightarrow (\text{CH}_3\text{COOH})_2$ .  
(ii) When a solute undergoes dissociation in solution then the number of solute particles increases. Hence, colligative property will be higher and the molar mass will be lower *e.g.*,  $\text{NaCl} \longrightarrow \text{Na}^+ + \text{Cl}^-$ .
- In a cubic crystal system, there are two types of voids known as octahedral and tetrahedral voids. If  $r$  is the radius of void and  $R$  is the radius of atom creating these voids then

$$\left(\frac{r}{R}\right)_{\text{octa}} = 0.414 \quad \text{and} \quad \left(\frac{r}{R}\right)_{\text{tetra}} = 0.225$$

The above radius ratio values indicate that octahedral void has larger radius hence, for maximum diameter of atom to be present in interstitial space.

$$r = 0.414 R$$

Also, in fcc,  $4R = \sqrt{2}a$

Diameter required  $(2r) = (2R) \times 0.414$

$$= \frac{a}{\sqrt{2}} \times 0.414 = \frac{400 \times 0.414}{\sqrt{2}} = 117 \text{ pm}$$

OR

- (i) When silicon is doped with boron which contains only three valence electrons, a hole is created at the place where the fourth valence electron is missing. This hole increases the conductivity and this type of semiconductors are called *p*-type semiconductors.
- (ii) As all the magnetic moments are aligned in one direction hence, the substance will show ferromagnetism.

9. Solubility of  $\text{H}_2\text{S}$  = 0.195 m = 0.195 mol in 1000 g of the solvent (water)

$$n_{\text{H}_2\text{O}} = \frac{1000 \text{ g}}{18 \text{ g mol}^{-1}} = 55.55 \text{ moles}$$

$$\therefore x_{\text{H}_2\text{S}} = \frac{n_{\text{H}_2\text{S}}}{n_{\text{H}_2\text{S}} + n_{\text{H}_2\text{O}}} = \frac{0.195}{0.195 + 55.55} = \frac{0.195}{55.745} = 0.0035$$

Pressure at STP = 0.987 bar

Applying Henry's law,  $p_{\text{H}_2\text{S}} = K_{\text{H}} \times x_{\text{H}_2\text{S}}$

$$\text{or } K_{\text{H}} = \frac{p_{\text{H}_2\text{S}}}{x_{\text{H}_2\text{S}}} = \frac{0.987 \text{ bar}}{0.0035} = 282 \text{ bar.}$$

10. Schottky defects exist in pairs to maintain electrical neutrality. So, equal number of cation and anion vacancies are generated.  
Ionic solids are doped with metal ions of higher valency and some cations of lower valency are displaced to maintain electrical neutrality. Hence, only cation vacancies are produced not anion vacancies.

11. (i) Given  $d = 8.55 \text{ g cm}^{-3}$ ,  $M = 93 \text{ g mol}^{-1}$ ,  $Z = 2$  (for bcc),  $N_A = 6.022 \times 10^{23}$ ,  $r = ?$

Using formula,

$$a^3 = \frac{M \times Z}{d \times N_A} = \frac{93 \times 2}{8.55 \times 6.022 \times 10^{23}}$$

$$\text{or, } a^3 = 3.61 \times 10^{-23} = 36.1 \times 10^{-24} \text{ cm}^3 \\ \therefore a = 3.304 \times 10^{-8} \text{ cm} = 330.4 \times 10^{-12} \text{ m} \\ = 330.4 \text{ pm}$$

For body-centred cubic,

$$r = \frac{\sqrt{3}}{4} a = 0.433 a = 0.433 \times 330.4 \text{ pm} \\ = 143.1 \text{ pm}$$

- (ii) The size and shape of a unit cell is determined by the lengths of the edges of the unit cell ( $a$ ,  $b$  and  $c$ ) which may or may not be mutually perpendicular and by the angles  $\alpha$ ,  $\beta$  and  $\gamma$  between the edges  $b$  and  $c$ ,  $c$  and  $a$  and  $a$  and  $b$  respectively.

12. Since, 1 atm =  $1.01325 \times 10^5$  pascal  
Hence, 2.5 atm =  $2.5 \times 1.01325 \times 10^5$  pascal  
=  $2.533125 \times 10^5$  pascal

By Henry's law  $p_{\text{CO}_2} = K_{\text{H}} x_{\text{CO}_2}$

$$x_{\text{CO}_2} = \frac{p_{\text{CO}_2}}{K_{\text{H}}} = \frac{2.533125 \times 10^5 \text{ Pa}}{1.67 \times 10^8 \text{ Pa}} \\ = 1.52 \times 10^{-3}$$

Also, 500 mL soda water (water containing  $\text{CO}_2$ )  
= 500 g (assuming density of water =  $1 \text{ g mL}^{-1}$ )

$$n_{\text{H}_2\text{O}} = \frac{500}{18} = 27.77 \text{ mol}$$

Let  $n_{\text{CO}_2} = n \text{ mol}$


$\therefore$  Total moles =  $27.77 + n$

$$\therefore x_{\text{CO}_2} = \frac{n}{27.77 + n} = 1.52 \times 10^{-3}$$

$$\therefore \frac{n}{27.77} = 1.52 \times 10^{-3} \quad (\because n \ll 27.77)$$

$$\therefore n_{\text{CO}_2} = 1.52 \times 10^{-3} \times 27.77 \text{ mol} \\ = 0.0422 \text{ mol} = 0.0422 \text{ mol} \times 44 \text{ g mol}^{-1} \\ = 1.8568 \text{ g}$$

13. (i) On heating ZnO loses oxygen according to the following reaction :



$\text{Zn}^{2+}$  ions are entrapped in the interstitial voids and electrons are entrapped in the neighbouring interstitial voids to maintain electrical neutrality. These electrons absorb visible light and radiate yellow colour.

- (ii) (a) 15 ppm means 15 parts per million, i.e., 15 parts of  $\text{CHCl}_3$  in  $10^6$  parts by mass of the solution.

$$\therefore \% \text{ by mass} = \frac{15}{10^6} \times 100 = 15 \times 10^{-4} \%$$

- (b) Taking 15 g chloroform in  $10^6$  g of the solution so, mass of solvent =  $10^6$  g



$$\text{Molar mass of CHCl}_3 = 12 + 1 + 3 \times 35.5 \\ = 119.5 \text{ g mol}^{-1}$$

$$\therefore n_{\text{CHCl}_3} = \frac{15}{119.5}$$

$$\therefore \text{Molality} = \frac{n_{\text{CHCl}_3}}{w_{\text{H}_2\text{O}}(\text{in g})} \times 1000 \\ = \frac{15/119.5}{10^6} \times 1000 = 1.25 \times 10^{-4} \text{ m}$$

14. This solution of benzene and toluene will boil at 25°C when the pressure above the solution is equal to the sum of the vapour pressures of benzene and toluene in the solution.

$$p_{\text{C}_6\text{H}_6}^\circ = 93.4 \text{ torr}, p_{\text{C}_6\text{H}_5\text{CH}_3}^\circ = 26.9 \text{ torr}$$

$$w_{\text{C}_6\text{H}_6} = 35 \text{ g}, w_{\text{C}_6\text{H}_5\text{CH}_3} = 65 \text{ g}$$

$$\text{Molar mass of C}_6\text{H}_6 = 12 \times 6 + 1 \times 6 \\ = 72 + 6 = 78 \text{ g mol}^{-1}$$

$$\text{Therefore, } n_{\text{C}_6\text{H}_6} = \frac{35}{78} = 0.449$$

$$\text{Molar mass of C}_6\text{H}_5\text{CH}_3 = 12 \times 7 + 8 \times 1 = 92 \text{ g mol}^{-1}$$

$$\text{Therefore, } n_{\text{C}_6\text{H}_5\text{CH}_3} = \frac{65}{92} = 0.707$$

Now, the mole fractions of C<sub>6</sub>H<sub>6</sub> and C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub> are

$$x_{\text{C}_6\text{H}_6} = \frac{n_{\text{C}_6\text{H}_6}}{n_{\text{C}_6\text{H}_6} + n_{\text{C}_6\text{H}_5\text{CH}_3}} \\ = \frac{0.449}{0.449 + 0.707} = \frac{0.449}{1.156} = 0.388$$

$$x_{\text{C}_6\text{H}_5\text{CH}_3} = \frac{n_{\text{C}_6\text{H}_5\text{CH}_3}}{n_{\text{C}_6\text{H}_5\text{CH}_3} + n_{\text{C}_6\text{H}_6}} \\ = \frac{0.707}{0.707 + 0.449} = \frac{0.707}{1.156} = 0.612$$

Therefore, vapour pressures of C<sub>6</sub>H<sub>6</sub> and C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub> are

$$p_{\text{C}_6\text{H}_6} = p_{\text{C}_6\text{H}_6}^\circ \times x_{\text{C}_6\text{H}_6} = 93.4 \times 0.388 = 36.24 \text{ torr}$$

$$p_{\text{C}_6\text{H}_5\text{CH}_3} = p_{\text{C}_6\text{H}_5\text{CH}_3}^\circ \times x_{\text{C}_6\text{H}_5\text{CH}_3} \\ = 26.9 \times 0.612 = 16.46 \text{ torr}$$

$\therefore$  Total vapour pressure of the solution,

$$P_{\text{total}} = p_{\text{C}_6\text{H}_6} + p_{\text{C}_6\text{H}_5\text{CH}_3} = 36.24 + 16.46 = 52.7 \text{ torr}$$

15. (i) This is because glass is a supercooled liquid (amorphous solid) and it flows down very slowly making the bottom portion slightly thicker.
- (ii) There is perfect arrangement of the constituent particles only at absolute zero (0 K). This arrangement corresponds to lowest energy. As temperature increases, deviation from perfectly ordered arrangement starts and number of defects increases with temperature. Hence, the defects are called thermodynamic defects.

- (iii) On applying high pressure, NaCl acquires the CsCl structure *i.e.*, the coordination number changes from 6 : 6 to 8 : 8.

OR

- (i) Given, edge length =  $a = 4.070 \text{ \AA}$   
Closest distance between two gold atoms in *fcc*

$$\text{lattice } (d) = 2r = \frac{2 \times a}{2\sqrt{2}}$$

$$\Rightarrow d = \frac{a}{\sqrt{2}} = \frac{4.070}{\sqrt{2}} = 2.878 \text{ \AA}$$

- (ii) Coordination number of each ion in *fcc* lattice is 12 hence, the number of nearest neighbours = 12

$$(iii) d = \frac{ZM}{N_A a^3} = \frac{4 \times 197}{6.02 \times 10^{23} \times (4.07 \times 10^{-8})^3} \\ = 19.4 \text{ g/cm}^3$$

16. (i) We know that

$$\pi V = n_2 RT$$

$$\pi V = \frac{w_2}{M_2} RT \text{ or } M_2 = \frac{w_2 RT}{\pi V}$$

$$\text{Here, } w_2 = 7 \text{ g}, \pi = 25 \text{ mm Hg} = \frac{25}{760} \text{ atm}$$

$$R = 0.08205 \text{ L atm mol}^{-1} \text{ deg}^{-1} \text{ and}$$

$$V = \frac{100}{1000} = 0.1 \text{ litre}$$

$$T = 310 \text{ K}$$

$$\therefore M_2 = \frac{7 \times 0.08205 \times 310 \times 760}{25 \times 0.1} \\ = 54126.7 \text{ g mol}^{-1}$$

- (ii) We know that,

$$\Delta T_b = \frac{1000 K_b w_2}{M_2 w_1 (\text{in g})} = \frac{1000 \times 1.72 \times 0.456}{152 \times 31.4} = 0.16$$


$$\therefore \Delta T_b = \text{Boiling point of solution} -$$

boiling point of solvent

$$\therefore 0.16 = \text{Boiling point of solution} - 56.30$$

$$\Rightarrow \text{Boiling point of solution} = 0.16 + 56.30 \\ = 56.46^\circ \text{C}$$

17. Calcium chloride ionises as follows :



$$\Rightarrow 1 \text{ mole of CaCl}_2 \text{ gives 2 moles of Cl}^- \text{ ions.}$$

*i.e.*,  $2 \times 6.02 \times 10^{23} \text{ Cl}^-$  ions are obtained from 1 mole of CaCl<sub>2</sub>.

$\therefore 3.01 \times 10^{22} \text{ Cl}^-$  ions will be obtained from

$$= \frac{3.01 \times 10^{22}}{2 \times 6.02 \times 10^{23}} = \frac{1}{40} \text{ mole of CaCl}_2$$

Thus, 200 mL solution contains  $\frac{1}{40}$  mole of CaCl<sub>2</sub>.

∴ 1000 mL solution will contain

$$= \frac{1}{40} \times \frac{1000}{200} = 0.125 \text{ mol of CaCl}_2$$

Therefore, molarity of solution is 0.125 M.

18. (i) In graphite, three electrons of each carbon atom are covalently bonded to three other C-atoms of its own layer. The fourth valence electron of each atom forms a delocalised layer of mobile electrons. This is why graphite is a conductor of electricity.

- (ii) In *hcp* lattice, each third layer is parallel to first layer *i.e.*, ABAB..... type arrangement. In *ccp* lattice, each fourth layer is parallel to the first layer *i.e.*, ABCABC..... type arrangement.

19. 
$$\begin{array}{ccccccc} & & \text{HA} & \rightleftharpoons & \text{H}^+ & + & \text{A}^- \\ \text{Initial} & & C \text{ mol L}^{-1} & & 0 & & 0 \\ \text{At equilibrium} & & (C - C\alpha) & & C\alpha & & C\alpha \end{array}$$
  
( $\alpha$  = degree of dissociation)

$$\Rightarrow [\text{H}^+] = C\alpha$$

$$\text{But } [\text{H}^+] = 10^{-2} \text{ M} \quad [\because \text{pH} = 2]$$

$$\therefore C\alpha = 10^{-2}$$

$$\text{or } \alpha = \frac{10^{-2}}{C} = \frac{10^{-2}}{0.1} = 0.1$$

Total number of particles after dissociation  
 $= C - C\alpha + C\alpha + C\alpha = C(1 + \alpha)$

$$\therefore \text{van't Hoff factor, } (i) = \frac{C(1 + \alpha)}{C}$$

$$= 1 + \alpha = 1 + 0.1 = 1.1$$

- ∴ Osmotic pressure ( $\pi$ ) of the monobasic acid  
 $= iCRT = 1.1 \times 0.1 \times 0.0821 \times 298$   
 $= 2.69 \text{ atm}$

20. Let 1 g mixture contains  $x$  g of  $\text{Na}_2\text{CO}_3$ .

$$\therefore \text{Number of moles of } \text{Na}_2\text{CO}_3 = \frac{x}{106} \text{ mol}$$

$$[\because \text{molar mass of } \text{Na}_2\text{CO}_3 = 106 \text{ g mol}^{-1}]$$

$$\text{and number of moles of } \text{NaHCO}_3 = \frac{1-x}{84}$$

$$[\because \text{molar mass of } \text{NaHCO}_3 = 84 \text{ g mol}^{-1}]$$



As the mixture contains equimolar amounts of the two therefore,

$$\frac{x}{106} = \frac{1-x}{84} \Rightarrow x = 0.558 \text{ g}$$

$$\therefore \text{No. of moles of } \text{Na}_2\text{CO}_3 = \frac{0.558}{106} = 0.00526$$

$$\text{and no. of moles of } \text{NaHCO}_3 = \frac{1-0.558}{84} = 0.00526 \text{ mol}$$

Reactions of HCl with  $\text{Na}_2\text{CO}_3$  and  $\text{NaHCO}_3$  take place as follows :



From the reaction, 1 mol of  $\text{Na}_2\text{CO}_3$  requires

2 moles of HCl

$$\therefore 0.00526 \text{ mol of } \text{Na}_2\text{CO}_3 \text{ will require } 2 \times 0.00526 = 0.01052 \text{ mole of HCl}$$

Similarly, 1 mol of  $\text{NaHCO}_3$  requires 1 mole of HCl

$$\therefore 0.00526 \text{ mol of } \text{NaHCO}_3 \text{ will require } 0.00526 \times 1 = 0.00526 \text{ mole of HCl}$$

$$\therefore \text{Total moles of HCl required} = 0.01052 + 0.00526 = 0.01578 \text{ mol}$$

0.1 M solution of HCl means that

0.1 mol of HCl is present in 1000 mL of solution

$$\therefore 0.01578 \text{ mol of HCl will be present in}$$

$$= \frac{1000 \times 0.01578}{0.1} = 157.8 \text{ mL of HCl solution.}$$

21. (i) For cubic closed packed structure,

$$\text{edge length, } a = \frac{4r}{\sqrt{2}} = 2\sqrt{2}r$$

$$= 2\sqrt{2} \times 125 \text{ pm} = 250\sqrt{2} \text{ pm} = 353.55 \text{ pm}$$

$$= 353.55 \times 10^{-12} \text{ m} = 353.55 \times 10^{-10} \text{ cm}$$


- (ii) Volume of 1 unit cell =  $(353.55 \times 10^{-10})^3 \text{ cm}^3$   
 $= 4.4193 \times 10^{-23} \text{ cm}^3$

$$4.4193 \times 10^{-23} \text{ cm}^3 = 1 \text{ unit cell}$$

$$\therefore 1 \text{ cm}^3 = \frac{1 \text{ unit cell}}{4.4193 \times 10^{-23}}$$

$$= 2.263 \times 10^{22} \text{ unit cells}$$

22. (i) Solubility in *n*-octane :



The order can be explained on the basis of 'like dissolves like'. *n*-Octane is a non polar solvent hence, non polar compounds will be highly soluble in *n*-octane. KCl is an ionic compound,  $\text{CH}_3\text{CN}$  is more polar than  $\text{CH}_3\text{OH}$  and cyclohexane is non polar.

Therefore, solubility increases with decreasing polarity.

- (ii) Applying the formula,

$$M_2 = \frac{1000 \times K_f \times w_2}{w_1 \times \Delta T_f}$$

$$M_{AB_2} = \frac{1000 \times 5.1 \times 1}{20 \times 2.3} = 110.87 \text{ g mol}^{-1}$$

$$M_{AB_4} = \frac{1000 \times 5.1 \times 1}{20 \times 1.3} = 196.15 \text{ g mol}^{-1}$$

Suppose atomic masses of A and B are 'a' and 'b' respectively. Then

Molar mass of  $AB_2 = a + 2b = 110.87 \text{ g mol}^{-1}$   
Molar mass of  $AB_4 = a + 4b = 196.15 \text{ g mol}^{-1}$   
Eqn. (ii) - Eqn. (i) gives  $2b = 85.28$  or  $b = 42.64$   
Substituting in eqn. (i) we get,  
 $a + 2 \times 42.64 = 110.87$  or  $a = 25.59$   
Thus, atomic mass of  $A = 25.59 \text{ u}$   
atomic mass of  $B = 42.64 \text{ u}$

23. (i) Curiosity to learn or discover new things and appropriate usage of resources are the values shown by Aamir.
- (ii) Semiconductors is the class of solids used in the components of electric devices.
- (iii) In electronic devices, there is need to control the current flow in the circuit which is possible only in semiconductors by appropriate doping. However, if conductors are used then such control is not possible and they conduct the whole amount of current from the power source.
- (iv) Conductivity range of semiconductors is  $10^{-6}$  to  $10^4 \text{ ohm}^{-1} \text{ m}^{-1}$ .
24. (i) KCl becomes violet coloured when heated in potassium vapours as there are excess of  $K^+$  ions which attract  $Cl^-$  ions and electrons are trapped at the site of anion forming F-centres. These electrons absorb energy from visible light and radiate complimentary violet colour.
- (ii) For cubic crystals,

$$d = \frac{Z \times M}{a^3 \times N_A} \quad \text{or,} \quad Z = \frac{d \times a^3 \times N_A}{M}$$

$$\therefore Z = \frac{(2.75 \text{ g cm}^{-3})(654 \times 10^{-10} \text{ cm})^3 \times (6.023 \times 10^{23})}{(39 + 80) \text{ g mol}^{-1}}$$

$$= 3.89 \approx 4$$

So, there are four formula units of KBr present per unit cell. Hence, it has face-centred cubic lattice.

- (iii) For the *bcc* structure, nearest neighbour distance ( $d$ ) is related to the edge length ( $a$ ) as

$$d = \frac{\sqrt{3}}{2} a$$

$$\text{or, } a = \frac{2}{\sqrt{3}} d = \frac{2}{1.732} \times 365.9 = 422.5 \text{ pm}$$

For *bcc* structure,  $Z = 2$

For sodium,  $M = 23$

$$\text{Density, } \rho = \frac{Z \times M}{a^3 \times N_A}$$

$$\rho = \frac{2 \times 23 \text{ g mol}^{-1}}{(422.5 \times 10^{-10} \text{ cm})^3 \times (6.02 \times 10^{23} \text{ mol}^{-1})}$$


$$= 1.013 \text{ g/cm}^3$$

OR

- (i) As the solid has same value of refractive index along all directions, it is isotropic in nature. It is because there is no long range order and the arrangement is irregular along all the directions and hence, amorphous. Being an amorphous solid, it would not show a clean cleavage when cut with a knife. Instead, it would break into pieces with irregular surfaces.

- (ii) (a) In solid state of sodium chloride ( $\text{NaCl}$ ), cations ( $\text{Na}^+$ ) and anions ( $\text{Cl}^-$ ) are held together by strong electrostatic (coulombic) forces hence, it does not carry electrical charges.

But in the presence of water, sodium chloride gets hydrolysed and splits in cations ( $\text{Na}^+$ ) and anions ( $\text{Cl}^-$ ) which are free to move and carry electrical charges through the aqueous medium.




Hence, sodium chloride which is insulator in solid state, carries charges in aqueous solution and becomes a good conductor.

- (b) The electronic configuration of  $\text{Na}^+$  is  $1s^2 2s^2 2p^6$  and that of  $\text{Cl}^-$  is  $1s^2 2s^2 2p^6 3s^2 3p^6$ . As no unpaired electron is present in both the ions hence, both will exhibit diamagnetic properties.
- (c) Schottky defect and metal excess defect are generally observed in sodium chloride.


25. (i) Camphor has very high molal depression constant ( $K_f = 39.70 \text{ K kg mol}^{-1}$ ) therefore, depression in freezing point will be large even when small quantity of a solute is added. This would be easily measurable even by an ordinary thermometer.

- (ii) Aquatic species need dissolved oxygen for breathing. As the solubility of gas, e.g.,  $\text{O}_2$  decreases with rise in temperature, less oxygen is available in warm water. Hence, they feel more comfortable in cold water when rather more dissolved oxygen is available.


# Concerned about your performance in **Class XII** Boards?




₹ 475



₹ 475



₹ 500



₹ 400

**Well, fear no more, help is at hand.....**

To excel, studying in right direction is more important than studying hard. Which is why we created the Excel Series. These books – for Physics, Chemistry, Biology & Mathematics – have been put together totally keeping in mind the prescribed syllabus and the pattern of CBSE's Board examinations, so that students prepare and practice with just the right study material to excel in board exams.

Did you know nearly all questions in CBSE's 2016 Board Examination were a part of our Excel books? That too fully solved !

## HIGHLIGHTS:

- Comprehensive theory strictly based on NCERT, complemented with illustrations, activities and solutions of NCERT questions
- Practice questions & Model Test Papers for Board Exams
- Value based questions
- Previous years' CBSE Board Examination Papers (Solved)
- CBSE Board Papers 2016 Included



Scan now with your smartphone or tablet

- (iii) Given,  $p_A^\circ = 450 \text{ mm}$ ,  $p_B^\circ = 700 \text{ mm}$ ,  
 $P_{\text{total}} = 600 \text{ mm}$

According to Raoult's law,  $p_A = x_A \times p_A^\circ$

$$p_B = x_B \times p_B^\circ = (1 - x_A) p_B^\circ$$

$$P_{\text{total}} = p_A + p_B = x_A p_A^\circ + (1 - x_A) p_B^\circ$$

$$= p_B^\circ + (p_A^\circ - p_B^\circ) x_A$$

On substituting the values, we get

$$600 = 700 + (450 - 700) x_A$$

$$\text{or } 250 x_A = 100$$

$$\text{or } x_A = \frac{100}{250} = 0.40$$

Thus, composition of the liquid mixture will be

mole fraction of A ( $x_A$ ) = 0.40,

mole fraction of B ( $x_B$ ) = 1 - 0.40 = 0.60

$$\therefore p_A = x_A \times p_A^\circ = 0.40 \times 450 \text{ mm} = 180 \text{ mm}$$

$$p_B = x_B \times p_B^\circ = 0.60 \times 700 \text{ mm} = 420 \text{ mm}$$


Thus, mole fraction of A in vapour phase

$$= \frac{p_A}{p_A + p_B} = \frac{180}{180 + 420} = 0.30$$

$$\text{Mole fraction of B in vapour phase} = 1 - 0.30 = 0.70$$

OR

- (i) This phenomenon is called endosmosis, i.e., movement of water inside the raisin which can be shown with the help of diagram as :



The process of osmosis is of immense biological as well as industrial importance. It is evident from the following examples :

- Movement of water from soil into plant roots and subsequently into upper portion of the plant is partly due to osmosis.
  - Preservation of meat against bacterial action by addition of salt.
  - Preservation of fruits against bacterial action by adding sugar. Bacteria in canned fruits lose water through the process of osmosis and become inactive.
- (ii) NaCl is a non-volatile solute hence, it lowers the vapour pressure of water and boiling point of water increases. Methyl alcohol is a volatile

solute hence, it increases the vapour pressure and results in decrease in boiling point of water.

26. (i) Alkali metal ions have large size and cannot fit into interstitial sites. Hence, Frenkel defect is not found in alkali metals.

- (ii) Radius of cation that just fits in the tetrahedral hole,

$$r = 0.225 R$$

$$\text{or, } r = 0.225 \times 195 = 43.875 \text{ pm}$$

For octahedral void,  $\frac{r}{R} = 0.414 \text{ to } 0.732$


$$\text{Radius ratio of given ions, } \frac{r_+}{r_-} = \frac{82}{195} = 0.4205$$

As the ratio lies in the range of radius of octahedral void hence, cation  $A^+$  can be slipped into the octahedral void of  $A^+B^-$  crystal.

- (iii) (a) Metal excess defects due to anionic vacancies result in the formation of *F*-centres. These *F*-centres produce colour in alkali metal halides.
- (b) When the electricity is produced by applying mechanical stress on some polar crystals, it is known as piezoelectric effect. Quartz shows this property.

OR

- (i) (a) According to band theory the energy gap between valence band and conduction band (Forbidden band) in insulator is very large whereas in a conductor valence band and conduction band overlap with each other, it has no forbidden band.
- (b) Conductors have no forbidden band whereas semiconductors have small forbidden band.



- (ii) Copper to oxygen ratio in  $\text{Cu}_2\text{O}$  is less than 2:1. Thus, the solid has  $\text{Cu}^+$  ions less than required. This solid has cation vacancies which are produced when some of  $\text{Cu}^+$  ions are replaced by  $\text{Cu}^{2+}$  ions and holes are produced. Thus, this substance is a *p*-type semiconductor.





# Kerala PET

- Choose the wrong statement in the following.
  - $\text{TiO}_2$  is used in the pigment industry.
  - $\text{MnO}_2$  is used in dry battery cells.
  - $\text{V}_2\text{O}_5$  catalyses the oxidation of  $\text{SO}_2$  in the manufacture of sulphuric acid.
  - Ziegler catalyst is used in the manufacture of high density polyethylene.
  - The 'silver' UK coins are made of Ag/Ni alloy.
- In aqueous solution,  $\text{Cr}^{2+}$  is stronger reducing agent than  $\text{Fe}^{2+}$ . This is because
  - $\text{Cr}^{2+}$  ion is more stable than  $\text{Fe}^{2+}$
  - $\text{Cr}^{3+}$  ion with  $d^3$  configuration has favourable crystal field stabilisation energy
  - $\text{Cr}^{3+}$  has half-filled configuration and hence more stable
  - $\text{Fe}^{3+}$  in aqueous solution is more stable than  $\text{Cr}^{3+}$
  - $\text{Fe}^{2+}$  ion with  $d^6$  configuration has favourable crystal field stabilisation energy.
- Choose the correct matching of transition metal ion and magnetic moment from the codes given below :  
(At. No : Ti = 22, V = 23, Fe = 26)  

Transition element	Magnetic moment (B.M.)
(A) Titanium (III)	(1) 4.9
(B) Vanadium (II)	(2) 1.73
(C) Iron (II)	(3) 3.87

  - (A) - (2), (B) - (3), (C) - (1)
  - (A) - (2), (B) - (1), (C) - (3)
  - (A) - (1), (B) - (2), (C) - (3)
  - (A) - (1), (B) - (3), (C) - (2)
  - (A) - (3), (B) - (2), (C) - (1)
- The standard enthalpy of formation of  $\text{H}_2\text{O}_{(l)}$  and  $\text{Fe}_2\text{O}_{3(s)}$  are respectively  $-286 \text{ kJ mol}^{-1}$  and  $-824 \text{ kJ mol}^{-1}$ . What is the standard enthalpy change for the following reaction?  

$$\text{Fe}_2\text{O}_{3(s)} + 3\text{H}_{2(g)} \longrightarrow 3\text{H}_2\text{O}_{(l)} + 2\text{Fe}_{(s)}$$
  - $-538 \text{ kJ mol}^{-1}$
  - $+538 \text{ kJ mol}^{-1}$
  - $-102 \text{ kJ mol}^{-1}$
  - $+34 \text{ kJ mol}^{-1}$
  - $-34 \text{ kJ mol}^{-1}$
- The correct descending order of the heat liberated (in kJ) during the neutralisation of the acids  $\text{CH}_3\text{COOH}$  (W), HF (X),  $\text{HCOOH}$  (Y) and HCN (Z) under identical conditions ( $K_a$  of  $\text{CH}_3\text{COOH} = 1.8 \times 10^{-5}$ ,  $\text{HCOOH} = 1.8 \times 10^{-4}$ ,  $\text{HCN} = 4.9 \times 10^{-10}$  and  $\text{HF} = 3.2 \times 10^{-4}$ ) is
  - $Y > X > Z > W$
  - $X > Y > W > Z$
  - $W > X > Y > Z$
  - $Z > W > Y > X$
  - $Z > Y > X > W$
- How many times a 0.1 M strong monobasic acid solution should be diluted so that pH of the resulting solution is tripled?  
 (a) 50 (b) 10 (c) 25 (d) 100 (e) 1000
- The equilibrium pressure for the reaction  $\text{MSO}_4 \cdot 2\text{H}_2\text{O}_{(s)} \rightleftharpoons \text{MSO}_{4(s)} + 2\text{H}_2\text{O}_{(g)}$  is  $\pi/4 \text{ atm}$  at 400 K. The  $K_p$  for the given reaction (in  $\text{atm}^2$ ) is
  - $\pi^2/4$
  - $\pi/6$
  - $\pi^2/16$
  - $\pi/16$
  - $16/\pi$
- Calculate the molality of a solution that contains 51.2 g of naphthalene, ( $\text{C}_{10}\text{H}_8$ ), in 500 mL of carbon tetrachloride. The density of  $\text{CCl}_4$  is 1.60 g/mL.
  - 0.250 m
  - 0.500 m
  - 0.750 m
  - 0.840 m
  - 1.69 m
- 31 g of ethylene glycol ( $\text{C}_2\text{H}_6\text{O}_2$ ) is mixed with 500 g of solvent ( $K_f$  of the solvent is  $2 \text{ K kg mol}^{-1}$ ). What is the freezing point of the solution in K? (freezing point of solvent = 273 K)
  - 272
  - 271
  - 270
  - 274
  - 275

10. The standard reduction potential for  $\text{Zn}^{2+}/\text{Zn}$ ,  $\text{Ni}^{2+}/\text{Ni}$  and  $\text{Fe}^{2+}/\text{Fe}$  are  $-0.76$ ,  $-0.23$  and  $-0.44$  V respectively. The reaction  $X + Y^{2+} \longrightarrow X^{2+} + Y$  will have more negative  $\Delta G$  value when X and Y are

- (a)  $X = \text{Ni}$ ;  $Y = \text{Fe}$       (b)  $X = \text{Ni}$ ;  $Y = \text{Zn}$   
 (c)  $X = \text{Fe}$ ;  $Y = \text{Zn}$       (d)  $X = \text{Zn}$ ;  $Y = \text{Ni}$   
 (e)  $X = \text{Fe}$ ;  $Y = \text{Ni}$

11. Thermal decomposition of ammonium dichromate gives

- (a)  $\text{N}_2$ ,  $\text{H}_2\text{O}$  and  $\text{Cr}_2\text{O}_3$       (b)  $\text{N}_2$ ,  $\text{NH}_3$  and  $\text{CrO}$   
 (c)  $(\text{NH}_4)_2\text{CrO}_4$  and  $\text{H}_2\text{O}$       (d)  $\text{N}_2$ ,  $\text{H}_2\text{O}$  and  $\text{CrO}_3$   
 (e)  $\text{N}_2$ ,  $\text{H}_2\text{O}$  and  $\text{CrO}$

12. An element crystallising in body centred cubic lattice has an edge length of 500 pm. If its density is  $4 \text{ g cm}^{-3}$ , the atomic mass of the element (in  $\text{g mol}^{-1}$ ) is (consider  $N_A = 6 \times 10^{23}$ )

- (a) 100      (b) 250  
 (c) 125      (d) 150  
 (e) 50

13. The rate constant of the reaction,  $2\text{N}_2\text{O}_5 \longrightarrow 4\text{NO}_2 + \text{O}_2$  at 300 K is  $3 \times 10^{-5} \text{ s}^{-1}$ . If the rate of the reaction at the same temperature is  $2.4 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$ , then the molar concentration of  $\text{N}_2\text{O}_5$  is

- (a) 0.4 M      (b) 0.8 M  
 (c) 0.04 M      (d) 0.08 M  
 (e) 0.6 M

14. In the reaction  $A \rightarrow \text{Products}$ , when the concentration of A was reduced from  $2.4 \times 10^{-2} \text{ M}$  to  $1.2 \times 10^{-2} \text{ M}$  the rate decreased 8 times at the same temperature. The order of the reaction is

- (a) 0      (b) 1  
 (c) 2      (d) 3  
 (e) 0.5

15. Enzymatic reactions are given in Column I and enzymes in Column II

Column I	Column II
(A) Maltose $\rightarrow$ Glucose	(i) Zymase
(B) Sucrose $\rightarrow$ Glucose + Fructose	(ii) Pepsin
(C) Glucose $\rightarrow$ Ethyl alcohol + $\text{CO}_2$	(iii) Maltase
(D) Starch $\rightarrow$ Maltose	(iv) Invertase
(E) Proteins $\rightarrow$ Amino acids	(v) Diastase

Choose the correct matching of enzymatic reaction and enzyme that catalyses the correct reaction from the codes given below :

- (a) (A) - (ii), (B) - (iv), (C) - (v), (D) - (iii), (E) - (i)  
 (b) (A) - (iii), (B) - (iv), (C) - (i), (D) - (v), (E) - (ii)  
 (c) (A) - (v), (B) - (iv), (C) - (ii), (D) - (i), (E) - (iii)  
 (d) (A) - (v), (B) - (iii), (C) - (iv), (D) - (ii), (E) - (i)  
 (e) (A) - (ii), (B) - (iii), (C) - (i), (D) - (v), (E) - (iv)

16. In which one of the following properties, physisorption and chemisorption resemble each other?

- (a) Force of attraction  
 (b) Enthalpy of adsorption  
 (c) Temperature effect  
 (d) Effect of surface area  
 (e) Number of adsorption layers

17. Consider the following two complex ions :  $[\text{CoF}_6]^{3-}$  and  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$ . Which of the following statement(s) is/are false?

- I. Both are octahedral.  
 II.  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$  is diamagnetic while  $[\text{CoF}_6]^{3-}$  is paramagnetic.  
 III. Both are outer orbital complexes.  
 IV. In both the complexes the central metal is in the same oxidation state.  
 (a) II and III      (b) II, III and IV  
 (c) III only      (d) III and IV  
 (e) I, II and IV

18. Chlorophyll is a coordination compound of

- (a) iron      (b) magnesium  
 (c) manganese      (d) chromium  
 (e) zinc.

19. Which one of the following is a benzenoid aromatic compound?

- (a) Furan      (b) Thiophene  
 (c) Pyridine      (d) Aniline  
 (e) Cyclopentadienyl anion

20. The major product obtained by the addition reaction of HBr to 4-methylpent-1-ene in the presence of peroxide is

- (a) 1-bromo-4-methylpentane  
 (b) 4-bromo-2-methylpentane  
 (c) 2-bromo-4-methylpentane  
 (d) 3-bromo-2-methylpentane  
 (e) 2-bromo-2-methylpentane.


21. Which one of the following involves nucleophilic addition?
- Kolbe's reaction of phenol
  - Williamson's synthesis of ethers
  - Reimer-Tiemann's reaction of phenol
  - Kolbe's electrolytic synthesis of ethane from sodium acetate
  - Aldol formation from ethanal
22. The number of possible stereoisomers of the compound  $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_3$  is
- 2
  - 4
  - 3
  - 5
  - 6
23. Some organic compounds are given in List I and their uses in List II. Choose the correct matching.
- | List I  | List II                     |
|---|-----------------------------|
| (A) Triiodomethane                                | (i) solvent for alkaloids   |
| (B) <i>p, p'</i> -Dichlorodiphenyltrichloroethane | (ii) propellant in aerosols |
| (C) Trichloromethane                              | (iii) antiseptic            |
| (D) Dichloromethane                               | (iv) insecticide            |
- (A) - (ii), (B) - (iv), (C) - (i), (D) - (iii)
  - (A) - (iii), (B) - (iv), (C) - (i), (D) - (ii)
  - (A) - (ii), (B) - (i), (C) - (iv), (D) - (iii)
  - (A) - (iii), (B) - (i), (C) - (iv), (D) - (ii)
  - (A) - (i), (B) - (iii), (C) - (ii), (D) - (iv)
24. The total number of monohalogenated products formed by halogenation of 2,4,4-trimethylhexane is
- 5
  - 7
  - 6
  - 8
  - 0
25. Number of acyclic structural isomers of the compound having the molecular formula  $\text{C}_4\text{H}_{10}\text{O}$  is
- 4
  - 5
  - 6
  - 7
  - 3
26. 375 mg of an alcohol reacts with required amount of methyl magnesium bromide and releases 140 mL of methane gas at STP. The alcohol is
- ethanol
  - n*-butanol
  - methanol
  - n*-propanol
  - phenol.
27. Predict the product (B) in the following sequence of reactions :
- $$\text{Ethylbenzene} \xrightarrow{\text{KMnO}_4 - \text{KOH}} \text{A} \xrightarrow{\text{H}_3\text{O}^+} \text{B}$$
- Benzaldehyde
  - Benzophenone
  - Benzene
  - Acetophenone
  - Benzoic acid
28. Freon 12 is manufactured from  $\text{CCl}_4$  by
- Wurtz reaction
  - Swarts reaction
  - Fittig reaction
  - Wurtz-Fittig reaction
  - Sandmeyer reaction.
29. Which one of the following can be prepared by Gabriel phthalimide synthesis?
- Aniline
  - o*-Toluidine
  - Benzylamine
  - N*-Methylethanamine
  - 4-Bromoaniline
30. 4-Nitrotoluene is treated with bromine to get compound 'P'. 'P' is reduced with Sn and HCl to get compound 'Q'. 'Q' is diazotised and the product is treated with phosphinic acid to get compound 'R'. 'R' is oxidized with alkaline  $\text{KMnO}_4$  to get compound 'S'. Compound 'S' is
- 2-bromo-4-hydroxybenzoic acid
  - benzoic acid
  - 4-bromobenzoic acid
  - 3-bromobenzoic acid
  - 2-bromobenzoic acid.
31. Narcotic analgesic is
- aspirin
  - paracetamol
  - codeine
  - zantac
  - cimetidine.
32. In double strand helix structure of DNA, heterocyclic base cytosine forms hydrogen bond with
- adenine
  - guanine
  - purine
  - thymine
  - uracil.
33. The amino acid containing mercaptan unit is
- leucine
  - glutamine
  - cysteine
  - lysine
  - isoleucine.
34. Which one of the following is a non-reducing sugar?
- Maltose
  - Lactose
  - Sucrose
  - Glucose
  - Fructose
35. In the hydrogen atomic spectrum, the emission of the least energetic photon takes place during the transition from  $n = 6$  energy level to  $n = \dots\dots\dots$  energy level.
- 1
  - 3
  - 5
  - 4
  - 2

36. If 27 g of water is formed during complete combustion of pure propene ( $C_3H_6$ ), the mass of propene burnt is  
(a) 42 g (b) 21 g (c) 14 g (d) 56 g (e) 40 g
37. When 2.46 g of a hydrated salt ( $MSO_4 \cdot xH_2O$ ) is completely dehydrated, 1.20 g of anhydrous salt is obtained. If the molecular weight of anhydrous salt is  $120 \text{ g mol}^{-1}$  what is the value of  $x$ ?  
(a) 2 (b) 4 (c) 5 (d) 6 (e) 7
38. Identify the T-shaped molecule in the following :  
(a)  $BF_3$  (b)  $NH_3$  (c)  $NF_3$  (d)  $ClF_3$  (e)  $PCl_3$
39. Which one of the following molecules has the least dipole moment?  
(a)  $H_2O$  (b)  $BeF_2$   
(c)  $NH_3$  (d)  $NF_3$   
(e)  $BF_3$
40. Dipole-dipole interaction energy between polar molecules in solids depends on the radius of the molecule ( $r$ ) and it is directly proportional to  
(a)  $(1/r^2)$  (b)  $(1/r^6)$   
(c)  $(1/r)$  (d)  $(1/r^5)$   
(e)  $(1/r^3)$
41. Critical density of a gas having molecular weight  $39 \text{ g mol}^{-1}$  is  $0.1 \text{ g cm}^{-3}$ . Its critical volume in  $\text{L mol}^{-1}$  is  
(a) 0.390 (b) 3.90  
(c) 0.039 (d) 39.0  
(e) 390
42. The various types of hydrides and examples of each type are given below :
- | Hydride type           | Compound      |
|------------------------|---------------|
| (A) Electron deficient | (i) LiH       |
| (B) Saline             | (ii) $CH_4$   |
| (C) Electron-precise   | (iii) $NH_3$  |
| (D) Interstitial       | (iv) $B_2H_6$ |
| (E) Electron rich      | (v) CrH       |
- Choose the correct matching from the codes given below :
- (a) (A) - (ii), (B) - (iv), (C) - (v), (D) - (iii), (E) - (i)  
(b) (A) - (iv), (B) - (i), (C) - (ii), (D) - (v), (E) - (iii)  
(c) (A) - (iv), (B) - (iii), (C) - (v), (D) - (ii), (E) - (i)  
(d) (A) - (v), (B) - (iii), (C) - (iv), (D) - (ii), (E) - (i)  
(e) (A) - (iv), (B) - (v), (C) - (i), (D) - (ii), (E) - (iii)
43. One mole of hydrazine ( $N_2H_4$ ) loses 10 moles of electrons in a reaction to form a new compound X. Assuming that all the nitrogen atoms in hydrazine appear in the new compound, what is the oxidation state of nitrogen in X?  
(Note : There is no change in the oxidation state of hydrogen in the reaction.)  
(a) -1 (b) -3 (c) +3 (d) +5 (e) +1
44. The low solubility of LiF and that of CsI in water are respectively due to which of the properties of the alkali metal ions?  
(a) Higher hydration enthalpy of  $Li^+$ , higher lattice enthalpy of  $Cs^+$ .  
(b) Smaller hydration enthalpy of  $Li^+$ , higher lattice enthalpy of  $Cs^+$ .  
(c) Smaller lattice enthalpy of  $Li^+$ , higher hydration enthalpy of  $Cs^+$ .  
(d) Smaller hydration enthalpy of  $Li^+$ , smaller lattice enthalpy of  $Cs^+$ .  
(e) Higher lattice enthalpy of  $Li^+$ , smaller hydration enthalpy of  $Cs^+$ .
45. The second ionization enthalpy of which of the following alkaline earth metals is the highest?  
(a) Ba (b) Mg (c) Ca (d) Sr (e) Be
46. Which one of the following group 16 elements does not exist in -2 oxidation state?  
(a) S (b) Se (c) O (d) Po (e) Te
47. In which one of the following compounds of xenon, highest number of lone pair of electrons is present on xenon?  
(a)  $XeF_6$  (b)  $XeF_4$   
(c)  $XeOF_4$  (d)  $XeO_3$   
(e)  $XeF_2$
48. The hybridized state of  $Al^{3+}$  in the complex ion formed when  $AlCl_3$  is treated with aqueous acid is  
(a)  $sp^3$  (b)  $dsp^2$  (c)  $sp^3d^2$  (d)  $sp^2d$  (e)  $sp^2$

## SOLUTIONS

- (e) : The silver UK coins are made of Cu-Ni alloy.
- (b) :  $Cr^{3+}$  ion with  $d^3$  ( $t_{2g}^3$ ) configuration has favourable crystal field stabilisation energy.
- (a) : Magnetic moment,  $\mu = \sqrt{n(n+2)}$  B.M.

Ion	Electronic Configuration	Number of unpaired electrons	$\mu$
$Ti^{3+}$	$d^1$	1	1.73
$V^{2+}$	$d^3$	3	3.87
$Fe^{2+}$	$d^6$	4	4.90



5. (b): From the  $K_a$  values, strength of acids is in the order :  
 $\text{HF} > \text{HCOOH} > \text{CH}_3\text{COOH} > \text{HCN}$   
 Stronger the acid, higher is the heat liberated during the neutralisation as lesser energy is used for ionisation.

$\therefore$  The correct order of heat liberated is  
 $X > Y > W > Z$

6. (d):  $\text{pH} = -\log [\text{H}^+]$   
 $\text{pH of } 0.1 \text{ M strong monobasic acidic solution} = 1$   
 $\text{pH of the resulting solution} = 3$

$$\therefore [\text{H}^+] = 10^{-3} = \frac{0.1}{100}$$

So, the original solution is diluted 100 times.

7. (c): The given reaction is :  
 $\text{MSO}_4 \cdot 2\text{H}_2\text{O}_{(s)} \rightleftharpoons \text{MSO}_{4(s)} + 2\text{H}_2\text{O}_{(g)}$   
 As the pressure is due to  $\text{H}_2\text{O}_{(g)}$  only hence,

$$K_p = (p_{\text{H}_2\text{O}})^2 = \left(\frac{\pi}{4}\right)^2 = \frac{\pi^2}{16} \text{ atm}^2$$

8. (b): Volume of  $\text{CCl}_4 = 500 \text{ mL}$   
 Density of  $\text{CCl}_4 = 1.60 \text{ g/mL}$   
 $\therefore \text{Mass of } \text{CCl}_4 (w_1) = 500 \times 1.60 = 800 \text{ g}$   
 Mass of Naphthalene ( $w_2$ ) = 51.2 g  
 Now,  $m = \frac{w_2 \times 1000}{M_2 \times w_1 (\text{in g})} = \frac{51.2 \times 1000}{128 \times 800} = 0.500 \text{ m}$

9. (b):  $\Delta T_f = K_f \times m$   
 $= K_f \times \frac{w_2 \times 1000}{M_2 \times w_1 (\text{in g})} = 2 \times \frac{31 \times 1000}{62 \times 500} = 2$   
 $\therefore \text{Freezing point of solution} = 273 - 2 = 271 \text{ K}$

10. (d):  $X + Y^{2+} \longrightarrow X^{2+} + Y$   
 $\Delta G = -nFE_{\text{cell}}$   
 Greater the value of  $E_{\text{cell}}$ , more negative will be the value of  $\Delta G$  and greater the difference between standard reduction potentials of  $X^{2+}$  and  $Y^{2+}$ , greater will be the value of  $E_{\text{cell}}$ .  
 Hence,  $X = \text{Zn}$  and  $Y = \text{Ni}$

11. (a):  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7 \xrightarrow{\Delta} \text{N}_2 + \text{Cr}_2\text{O}_3 + 4\text{H}_2\text{O}$

12. (d): For bcc,  $Z = 2$

$$d = \frac{Z \times M}{N_A \times a^3}$$

$$\Rightarrow M = \frac{d \times N_A \times a^3}{Z}$$

$$\therefore M = \frac{4 \times 6 \times 10^{23} \times (500 \times 10^{-10})^3}{2}$$

$$= \frac{4 \times 6 \times 10^{23} \times 125 \times 10^{-24}}{2} = 150 \text{ g mol}^{-1}$$

13. (b): Rate constant,  $k = 3 \times 10^{-5} \text{ s}^{-1}$

From the unit of  $k$  it is clear that the reaction is of first order.

$$\Rightarrow \text{Rate} = k[\text{N}_2\text{O}_5]$$

$$\Rightarrow [\text{N}_2\text{O}_5] = \frac{\text{Rate}}{k} = \frac{2.4 \times 10^{-5}}{3 \times 10^{-5}} \text{ mol dm}^{-3}$$

$$= 0.8 \text{ mol dm}^{-3}$$

14. (d):  $\frac{r_1}{r_2} = \left[ \frac{2.4 \times 10^{-2}}{1.2 \times 10^{-2}} \right]^x$   
 $8 = (2)^x \Rightarrow (2)^3 = (2)^x$   
 $\therefore x = 3$

Hence, the order of reaction is 3.

15. (b): Maltose  $\xrightarrow{\text{Maltase}}$  Glucose

Sucrose  $\xrightarrow{\text{Invertase}}$  Glucose + Fructose



Glucose  $\xrightarrow{\text{Zymase}}$  Ethyl alcohol +  $\text{CO}_2$

Starch  $\xrightarrow{\text{Diastase}}$  Maltose

Proteins  $\xrightarrow{\text{Pepsin}}$  Amino acids

16. (d): Physisorption and chemisorption both increase with increase in surface area.

17. (c):  $[\text{CoF}_6]^{3-}$ ,




Hence, due to presence of 4 unpaired electrons  $[\text{CoF}_6]^{3-}$  is paramagnetic and  $\text{F}^-$  being a weak field ligand forms an outer orbital complex.

$[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$ ,




$[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$ :






Hence,  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$  is diamagnetic and  $\text{C}_2\text{O}_4^{2-}$  causes pairing of electrons and forms an inner orbital complex.

18. (b)



Benzenoid aromatic compounds have a benzene ring intact.



21. (e): Kolbe's reaction of phenol: It is an electrophilic substitution reaction.


Williamson's synthesis of ethers: It occurs by  $\text{S}_{\text{N}}2$  mechanism.

Reimer-Tiemann's reaction of phenol: It is an example of electrophilic substitution reaction involving dichlorocarbene as the electrophile.

Kolbe's electrolytic synthesis of ethane from sodium acetate: It involves free radical mechanism.

Aldol formation from ethanal: It is nucleophilic addition reaction.

22. (a):  $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_3$  will show geometrical isomerism.




23. (b): Triiodomethane – antiseptic

*p, p'*-Dichlorodiphenyltrichloroethane – insecticide



Trichloromethane – solvent for alkaloids

Dichloromethane – propellant in aerosols.






Six structural isomers corresponding to the substitution of 6 types of H atoms are possible.

25. (d): Four alcohols are possible:



and three ethers are possible:




$$\therefore 140 \text{ mL of } \text{CH}_4 \equiv \frac{1}{22400} \times 140 = 6.25 \times 10^{-3} \text{ mol of alcohol}$$

$$\text{Now, } n = \frac{w}{M}$$


$$\Rightarrow \text{Molecular mass of alcohol} = \frac{w}{n} = \frac{375 \times 10^{-3}}{6.25 \times 10^{-3}} = 60 \text{ g mol}^{-1}$$

Hence, the alcohol is *n*-propanol ( $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ ).

27. (e):



28. (b): Freon 12 is manufactured from  $\text{CCl}_4$  by Swarts reaction.



29. (c): Only primary aliphatic amines can be prepared by Gabriel phthalimide synthesis.

Hence, benzylamine can be prepared from Gabriel phthalimide synthesis.

The reaction scheme shows the synthesis of 2-bromobenzoic acid from 4-nitrotoluene. 
 1. 4-Nitrotoluene (a benzene ring with a methyl group at position 1 and a nitro group at position 4) reacts with  $\text{Br}_2$  to form (P), 4-bromo-4-nitrotoluene (a benzene ring with a methyl group at position 1, a nitro group at position 4, and a bromine atom at position 4).
 2. (P) reacts with  $\text{Sn/HCl}$  to form (Q), 4-bromoaniline (a benzene ring with a methyl group at position 1, an amino group at position 4, and a bromine atom at position 4).
 3. (Q) is treated with (i)  $\text{NaNO}_2/\text{HCl}$  and (ii)  $\text{H}_3\text{PO}_2$  to form (R), 2-bromo-1-methylbenzene (a benzene ring with a methyl group at position 1 and a bromine atom at position 2).
 4. (R) is oxidized with  $\text{Alk. KMnO}_4$  to form 2-bromobenzoic acid (a benzene ring with a carboxylic acid group at position 1 and a bromine atom at position 2).

**32. (b):** Cytosine forms three hydrogen bonds with guanine.

**34. (c):** Sucrose is a disaccharide in which the two monosaccharide units are linked through their reducing centres.

$$E_2 - E_1 > E_3 - E_2 > E_4 - E_3 \dots\dots$$

Hence, the emission of least energetic photon takes place during the transition from  $n = 6$  energy level to  $n = 5$  energy level.

54 g of  $\text{H}_2\text{O} \equiv 42 \text{ g of propene}$

$$\therefore 27 \text{ g of H}_2\text{O} = \frac{42}{54} \times 27 = 21 \text{ g}$$

$$\text{Mass of water} = 2.46 - 1.20 = 1.26 \text{ g}$$

$$\therefore \text{No. of moles of H}_2\text{O} = \frac{1.26}{18} = 0.07$$

$$\text{No. of moles of } \text{MSO}_4 = \frac{1.20}{120} = 0.01$$

$\therefore$  0.07 moles of water associated with 0.01 mole of salt.

Thus, there are 7 moles of water per mole of anhydrous salt.

$\text{NF}_3$  —  $sp^3$  hybridisation, pyramidal  
 $\text{ClF}_3$  —  $sp^3d$  hybridisation, T-shaped  
 $\text{PCl}_3$  —  $sp^3$  hybridisation, pyramidal

**39. (b, e) :**  $\text{BeF}_2$  has linear shape and  $\text{BF}_3$  is symmetrical hence, both have zero dipole moment.

**40. (e):** Dipole-dipole interaction energy between stationary polar molecules (as present in the solids) is proportional to  $1/r^3$ .

**41. (a):** Critical density =  $0.1 \text{ g cm}^{-3} = 100 \text{ g L}^{-1}$

$$\begin{aligned}\text{Critical volume} &= \frac{\text{Molecular weight}}{\text{Critical density}} \\ &= \frac{39}{100} = 0.39 \text{ L mol}^{-1}\end{aligned}$$

**42. (b):** Electron deficient –  $\text{B}_2\text{H}_6$

Saline – LiH

Electron precise – CH<sub>4</sub>

Interstitial – CrH

Electron rich – NH<sub>3</sub>

43. (c) :  $\text{N}_2\text{H}_4 \longrightarrow \text{X}$

No. of electrons lost per N-atom = 5

Oxidation state of N in new compound =  $-2 + 5 = +3$

44. (e)

**45. (e) :** On moving down the group, ionization energy decreases. This is due to the fact that the electrons in lower shells experience higher force of attraction from the nucleus. Thus, it is difficult for them to ionise.

46. (d)

47. (e):  $\text{XeF}_2$ : 3 lone pairs and 2 bond pairs.

**48. (c):**  $\text{AlCl}_3$  forms  $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$  complex when treated with aqueous acid and hybridisation of Al is  $sp^3 d^2$  in this complex.

**Buy MTG Books Online from [www.mtg.in](http://www.mtg.in)**

get

**25%** on orders above ₹ 999/-  
Apply coupon code - **COUPON25**  
**OFF\***

**MTG Books are also available on**

\*offer not applicable on combos

[www.flipkart.com](http://www.flipkart.com) | [www.amazon.in](http://www.amazon.in) | [www.snapdeal.com](http://www.snapdeal.com)

- The half-life period of a 1<sup>st</sup> order reaction is 60 minutes. What percentage will be left over after 240 minutes?  
(a) 6.25% (b) 4.25%  
(c) 5% (d) 6%
- Which of the following is not a colligative property?  
(a) Osmotic pressure  
(b) Optical activity  
(c) Depression in freezing point  
(d) Elevation in boiling point
- The contribution of particle at the edge centre of a particular unit cell is,  
(a)  $\frac{1}{2}$  (b)  $\frac{1}{4}$   
(c) 1 (d)  $\frac{1}{8}$
- When an electrolyte is dissociated in solution, the van't Hoff factor ( $i$ ) is  
(a)  $> 1$  (b)  $< 1$  (c)  $= 0$  (d)  $= 1$
- Which of the following is incorrect in a galvanic cell?  
(a) Oxidation occurs at anode.  
(b) Reduction occurs at cathode.  
(c) The electrode at which electrons are gained is called cathode.  
(d) The electrode at which electrons are lost is called cathode.
- A secondary cell is one which  
(a) can be recharged  
(b) can be recharged by passing current through it in the same direction  
(c) can be recharged by passing current through it in the opposite direction  
(d) cannot be recharged.
- Osmotic pressure of the solution can be increased by  
(a) increasing the temperature of the solution  
(b) decreasing the temperature of the solution  
(c) increasing the volume of the vessel  
(d) diluting the solution.
- The amount of current in faraday required for the reduction of 1 mol of  $\text{Cr}_2\text{O}_7^{2-}$  ions to  $\text{Cr}^{3+}$  is  
(a) 1 F (b) 2 F (c) 6 F (d) 4 F
- For a chemical reaction,  
 $m\text{A} \rightarrow x\text{B}$ , the rate law is  $r = k[\text{A}]^2$ .  
If the concentration of A is doubled, the reaction rate will be  
(a) doubled (b) quadrupled  
(c) increased by 8 times (d) unchanged.
- Schottky defect in a crystal is observed when  
(a) unequal number of cations and anions are missing from the lattice  
(b) equal number of cations and anions are missing from the lattice  
(c) an ion leaves its normal site and occupies an interstitial site  
(d) no ion is missing from its lattice site.
- $3\text{A} \longrightarrow 2\text{B}$ , rate of reaction,  $+\frac{d[\text{B}]}{dt}$  is equal to  
(a)  $-\frac{3}{2}\frac{d[\text{A}]}{dt}$  (b)  $-\frac{2}{3}\frac{d[\text{A}]}{dt}$   
(c)  $+2\frac{d[\text{A}]}{dt}$  (d)  $-\frac{1}{3}\frac{d[\text{A}]}{dt}$
- The activation energy of a chemical reaction can be determined by  
(a) evaluating rate constants at two different temperatures  
(b) changing the concentration of reactants

- (c) evaluating the concentration of reactants at two different temperatures  
(d) evaluating rate constant at standard temperature.

13. Which of the following statements is incorrect *w.r.t.* physisorption?

- (a) The forces involved are van der Waals' forces.  
(b) More easily liquifiable gases are adsorbed easily.  
(c) Under high pressure it results into multi-molecular layer on adsorbent surface.  
(d)  $\Delta H_{\text{adsorption}}$  is low and +ve.


14. Sulphur sol contains

- (a) discrete S-atoms  
(b) discrete S-molecules  
(c) large aggregates of S-molecules  
(d) water dispersed in solid sulphur.

15. Reactions in zeolite catalyst depend on

- (a) pores (b) apertures  
(c) size of cavity (d) all of these.

16. IUPAC name of the compound




is

- (a) 1-bromobut-2-ene (b) 2-bromo-2-butene  
(c) bromobutene (d) 1-bromobut-3-ene.

17. Replacement of Cl of Chlorobenzene to give phenol requires drastic conditions, but Cl of 2,4-dinitrochlorobenzene is readily replaced. This is because


- (a)  $-\text{NO}_2$  group makes the ring electron rich at *ortho* and *para* positions  
(b)  $-\text{NO}_2$  group withdraws electrons from *meta* position  
(c)  $-\text{NO}_2$  donates electrons at *meta* position  
(d)  $-\text{NO}_2$  withdraws electrons from *ortho* and *para* positions.

18. In the reaction :



- (a)  $\text{C}_2\text{H}_4$  (b)  $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$   
(c)  $\text{CH}_3\text{CH}_2\text{OSO}_3\text{H}$  (d)

19. Which of the following compounds is most acidic?



20. Benzene carbaldehyde is reacted with concentrated NaOH solution to give the products A and B. The product A can be used as food preservative and the product B is an aromatic hydroxy compound where OH group is linked to  $sp^3$  hybridised carbon atom next to benzene ring. The products A and B are respectively,

- (a) sodium benzoate and phenol  
(b) sodium benzoate and phenyl methanol  
(c) sodium benzoate and cresol  
(d) sodium benzoate and picric acid.

21. The reaction which involves dichlorocarbene as an electrophile is

- (a) Reimer-Tiemann reaction  
(b) Kolbe's reaction  
(c) Friedel-Crafts' acylation  
(d) Fittig's reaction.


22. Ethanol is converted into ethoxyethane

- (a) by heating excess of ethanol with conc.  $\text{H}_2\text{SO}_4$  at  $140^\circ\text{C}$   
(b) by heating ethanol with excess of conc.  $\text{H}_2\text{SO}_4$  at  $443\text{ K}$   
(c) by treating with conc.  $\text{H}_2\text{SO}_4$  at room temperature  
(d) by treating with conc.  $\text{H}_2\text{SO}_4$  at  $273\text{ K}$ .

23. An organic compound X is oxidised by using acidified  $\text{K}_2\text{Cr}_2\text{O}_7$  solution. The product obtained reacts with phenyl hydrazine but does not answer silver mirror test. The compound X is

- (a) 2-propanol (b) ethanal  
(c) ethanol (d)  $\text{CH}_3\text{CH}_2\text{CH}_3$

24. Predict the product 'C' in the following series of reactions :



- (a) (b)  $\text{CH}_3\text{CH}(\text{OH})\text{C}_6\text{H}_5$   
(c)  $\text{CH}_3\text{CH}(\text{OH})\text{C}_2\text{H}_5$  (d)  $(\text{CH}_3)_2\text{C}(\text{OH})\text{C}_6\text{H}_5$

25. The number of oxygen atoms in 4.4 g of  $\text{CO}_2$  is

- (a)  $1.2 \times 10^{23}$  (b)  $6 \times 10^{22}$   
(c)  $6 \times 10^{23}$  (d)  $12 \times 10^{23}$

26. If the bond energies of H—H, Br—Br and H—Br are 433, 192 and 364 kJ mol<sup>-1</sup> respectively, then  $\Delta H^\circ$  for the reaction :
- $$\text{H}_{2(g)} + \text{Br}_{2(g)} \longrightarrow 2\text{HBr}_{(g)}$$
- (a) - 261 kJ (b) + 103 kJ  
(c) + 261 kJ (d) - 103 kJ
27. In the reaction;  $\text{Fe}(\text{OH})_{3(s)} \rightleftharpoons \text{Fe}^{3+}_{(aq)} + 3\text{OH}^{-}_{(aq)}$ , if the concentration of OH<sup>-</sup> ions is decreased by  $\frac{1}{4}$  times, then the equilibrium concentration of Fe<sup>3+</sup> will increase by
- (a) 8 times (b) 16 times  
(c) 64 times (d) 4 times.
28. The correct statement regarding entropy is
- (a) at absolute zero temperature, entropy of a perfectly crystalline solid is zero  
(b) at absolute zero temperature, the entropy of a perfectly crystalline substance is +ve  
(c) at absolute zero temperature, the entropy of all crystalline substances is zero  
(d) at 0°C, the entropy of a perfect crystalline solid is zero.
29. Equilibrium constants  $K_1$  and  $K_2$  for the following equilibria
- $$(1) \text{NO}_{(g)} + \frac{1}{2} \text{O}_{2(g)} \rightleftharpoons \text{NO}_{2(g)}$$
- $$(2) 2\text{NO}_{2(g)} \rightleftharpoons 2\text{NO}_{(g)} + \text{O}_{2(g)}$$
- are related as :
- (a)  $K_1 = \sqrt{K_2}$  (b)  $K_2 = \frac{1}{K_1}$   
(c)  $K_1 = 2K_2$  (d)  $K_2 = \frac{1}{K_1^2}$
30. van-Arkel method of refining zirconium involves
- (a) removing all oxygen and nitrogen impurities  
(b) removing CO impurity  
(c) removing hydrogen impurity  
(d) removing silica impurity.
31. The composition of 'copper matte' is
- (a) Cu<sub>2</sub>S + FeS (b) Cu<sub>2</sub>S + Cu<sub>2</sub>O  
(c) Cu<sub>2</sub>S + FeO (d) Cu<sub>2</sub>O + FeS
32. The complex formed when Al<sub>2</sub>O<sub>3</sub> is leached from bauxite using concentrated NaOH solution is
- (a) Na[Al(OH)<sub>4</sub>] (b) NaAl<sub>2</sub>O<sub>4</sub>  
(c) Na<sub>2</sub>[Al(OH)<sub>3</sub>] (d) Na<sub>2</sub>AlO<sub>2</sub>
33. The property which is not true about fluorine is
- (a) most of its reactions are exothermic  
(b) it forms only one oxo acid  
(c) highest electronegativity  
(d) high F-F bond dissociation enthalpy.
34. Which is true regarding nitrogen?
- (a) Less electronegative  
(b) Has low ionisation enthalpy  
(c) *d*-orbitals are available  
(d) Ability to form *pπ-pπ* bonds with itself
35. The shape of XeF<sub>6</sub> is
- (a) square planar (b) distorted octahedral  
(c) square pyramidal (d) pyramidal.
36. The number of isomers possible for the octahedral complex  $[\text{CoCl}_2(en)(\text{NH}_3)_2]^+$  is
- (a) two (b) three  
(c) no isomer (d) four isomers.
37. CO is a stronger ligand than Cl<sup>-</sup>, because
- (a) CO is a neutral molecule.  
(b) CO has  $\pi$ -bonds.  
(c) CO is poisonous.  
(d) CO is more reactive.
38. The bivalent metal ion having maximum paramagnetic behaviour among the first transition series elements is
- (a) Mn<sup>2+</sup> (b) Cu<sup>2+</sup>  
(c) Sc<sup>2+</sup> (d) Cu<sup>+</sup>
39. When a brown compound of Mn (A) is treated with HCl, it gives a gas (B). The gas (B) taken in excess reacts with NH<sub>3</sub> to give an explosive compound (C).  
The compounds A, B and C are
- (a) A = MnO<sub>2</sub>, B = Cl<sub>2</sub>, C = NCl<sub>3</sub>  
(b) A = MnO, B = Cl<sub>2</sub>, C = NH<sub>3</sub>Cl  
(c) A = Mn<sub>3</sub>O<sub>4</sub>, B = Cl<sub>2</sub>, C = NCl<sub>3</sub>  
(d) A = MnO<sub>3</sub>, B = Cl<sub>2</sub>, C = NCl<sub>2</sub>
40. Mn<sup>2+</sup> compounds are more stable than Fe<sup>2+</sup> compounds towards oxidation to their +3 state, because
- (a) Mn<sup>2+</sup> is more stable with high 3<sup>rd</sup> ionisation energy  
(b) Mn<sup>2+</sup> is bigger in size  
(c) Mn<sup>2+</sup> has completely filled *d*-orbitals  
(d) Mn<sup>2+</sup> does not exist.



41. Which of the following sequence is correct regarding field strength of ligands as per spectrochemical series?
- $\text{SCN}^- < \text{F}^- < \text{CN}^- < \text{CO}$
  - $\text{F}^- < \text{SCN}^- < \text{CN}^- < \text{CO}$
  - $\text{CN}^- < \text{F}^- < \text{CO} < \text{SCN}^-$
  - $\text{SCN}^- < \text{CO} < \text{F}^- < \text{CN}^-$
42. As per IUPAC norms, the name of the complex  $[\text{Co}(\text{en})_2(\text{ONO})\text{Cl}]\text{Cl}$  is
- Chloridobis(ethane-1,2-diamine)nitro-O-cobalt (III) chloride.
  - Chlorobis(ethylenediamine)nitro-O-cobalt (III) chloride.
  - Chloridodi(ethylene diamine)nitrocobalt (III) chloride.
  - Chloroethylenediaminenitro-O-cobalt (III) chloride.
43. In the following sequence of reactions;
- $$\text{A} \xrightarrow{\text{Reduction}} \text{B} \xrightarrow{\text{HNO}_2} \text{CH}_3\text{CH}_2\text{OH}$$
- The compound A is
- propane nitrile
  - ethane nitrile
  - nitromethane
  - methyl isocyanate.
44. An organic compound A on reduction gives compound B, which on reaction with trichloro methane and caustic potash forms C. The compound 'C' on catalytic reduction gives N-methyl benzenamine, the compound 'A' is
- nitrobenzene
  - nitromethane
  - methanamine
  - benzenamine.
45. Which of the following gives positive Fehling's solution test?
- Sucrose
  - Glucose
  - Fats
  - Protein
46. A liquid can exist only
- between triple point and critical point
  - at any temperature above melting point
  - between melting point and critical point
  - between boiling and melting points.
47. The energy of electron in the  $n^{\text{th}}$  Bohr orbit of H-atom is
- $\frac{-13.6}{n^2} \text{eV}$
  - $\frac{-13.6}{n} \text{eV}$
  - $\frac{-13.6}{n^4} \text{eV}$
  - $\frac{-13.6}{n^3} \text{eV}$
48. Consider the following sets of quantum numbers : Which of the below setting is not permissible arrangement of electrons in an atom?
- |     | $n$ | $l$ | $m$ | $s$            |
|-----|-----|-----|-----|----------------|
| (a) | 4   | 0   | 0   | $-\frac{1}{2}$ |
| (b) | 5   | 3   | 0   | $+\frac{1}{2}$ |
| (c) | 3   | 2   | -2  | $-\frac{1}{2}$ |
| (d) | 3   | 2   | -3  | $+\frac{1}{2}$ |
49. The increasing order of bond orders of  $\text{O}_2$ ,  $\text{O}_2^+$ ,  $\text{O}_2^-$  and  $\text{O}_2^{2-}$  is
- $\text{O}_2^+, \text{O}_2, \text{O}_2^-, \text{O}_2^{2-}$
  - $\text{O}_2^-, \text{O}_2, \text{O}_2^+, \text{O}_2$
  - $\text{O}_2, \text{O}_2^+, \text{O}_2^-, \text{O}_2^{2-}$
  - $\text{O}_2^-, \text{O}_2, \text{O}_2^+, \text{O}_2^{2-}$
50. HCl gas is covalent and NaCl is an ionic compound. This is because
- sodium is highly electropositive
  - hydrogen is a non-metal
  - HCl is a gas
  - electronegativity difference between H and Cl is less than 2.1.
51. Which of the following is not true?
- In vulcanisation the rubber becomes harder and stronger.
  - Natural rubber has 'trans' configuration at every double bond.
  - Buna-S is a co-polymer of butene and styrene.
  - Natural rubber is 1,4-polymer of isoprene.
52. Which of the following is a polyamide?
- Nylon-6, 6
  - Terylene
  - Polythene
  - Buna-S
53. Which of the following is correct about H-bonding in DNA?
- A — T, G — C
  - A — G, T — G
  - G — T, A — C
  - A — A, T — T
54. Which of the following is employed as tranquilizer?
- Equanil
  - Naproxen
  - Tetracyclin
  - Dettol
55. Reactivity order of halides for dehydrohalogenation is
- $\text{R} - \text{F} > \text{R} - \text{Cl} > \text{R} - \text{Br} > \text{R} - \text{I}$
  - $\text{R} - \text{I} > \text{R} - \text{Br} > \text{R} - \text{Cl} > \text{R} - \text{F}$
  - $\text{R} - \text{I} > \text{R} - \text{Cl} > \text{R} - \text{Br} > \text{R} - \text{F}$
  - $\text{R} - \text{F} > \text{R} - \text{I} > \text{R} - \text{Br} > \text{R} - \text{Cl}$

56. Main axis of diatomic molecule is  $z$ . The orbitals  $p_x$  and  $p_y$  overlap to form  
 (a)  $\pi$ -molecular orbital (b)  $\sigma$ -molecular orbital  
 (c)  $\delta$ -molecular orbital (d) no bond is formed.
57. The hybridisation of C in diamond, graphite and ethyne is in the order  
 (a)  $sp^3, sp, sp^2$  (b)  $sp^3, sp^2, sp$   
 (c)  $sp, sp^2, sp^3$  (d)  $sp^2, sp^3, sp$
58. A miscible mixture of  $C_6H_6 + CHCl_3$  can be separated by  
 (a) sublimation (b) distillation  
 (c) filtration (d) crystallisation.
59. An organic compound contains C = 40%, H = 13.33% and N = 46.67%. Its empirical formula is  
 (a)  $C_2H_2N$  (b)  $C_3H_7N$   
 (c)  $CH_4N$  (d) CHN
60. Electrophile that participates in nitration of benzene is  
 (a)  $NO^+$  (b)  $NO_2^+$  (c) NO (d)  $NO_3^-$

### SOLUTIONS

1. (a):  $t_{1/2} = \frac{0.693}{k} \Rightarrow \frac{0.693}{t_{1/2}} = k \Rightarrow \frac{0.693}{60} = k$

$$k = 0.01155 \text{ min}^{-1}$$

$$k = \frac{2.303}{t} \log \left( \frac{a}{a-x} \right)$$

Let the initial amount ( $a$ ) be 100.

$$0.01155 \text{ min}^{-1} = \frac{2.303}{240 \text{ min}} \log \left( \frac{100}{a-x} \right)$$

$$\frac{0.01155 \text{ min}^{-1} \times 240 \text{ min}}{2.303} = \log \left( \frac{100}{a-x} \right)$$

$$1.204 = \log 100 - \log (a-x)$$

$$1.204 = 2 - \log (a-x)$$

$$\log (a-x) = 2 - 1.204$$

$$\log (a-x) = 0.796$$


$$(a-x) = 6.25\%$$

2. (b): Colligative property depends on the concentration of solution and not on the type of particles, while optical activity depends on the type of particles.
3. (b): Each particle at edge centre is surrounded by four unit cells thus, its contribution to each unit cell is  $1/4$ .
4. (a): van't Hoff factor ( $i$ ) depends on the number of particles of solute. On dissociation of an electrolyte, number of particles of solute increases, therefore  $i > 1$ .

5. (d): The electrode at which electrons are lost is called anode.

6. (c)

7. (a):  $\pi = CRT = \frac{n}{V} RT$   
 $\pi \propto T$



9. (b):  $r = k[A]^2$

When concentration of A is doubled,

$$r = k[2A]^2$$

$$r = 4k[A]^2$$

10. (b)

11. (b):  $-\frac{1}{3} \frac{d[A]}{dt} = +\frac{1}{2} \frac{d[B]}{dt}$

$$-\frac{2}{3} \frac{d[A]}{dt} = +\frac{d[B]}{dt}$$


12. (a):  $\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$

13. (d):  $\Delta H_{\text{adsorption}}$  is always negative.


14. (c): Sulphur sol consists of particles containing a thousand or more of  $S_8$  sulphur molecules.

15. (d) 16. (a)


17. (d):  $NO_2$  withdraws electrons from the carbon bearing chlorine group making the C—Cl bond weak.



19. (c): Due to electron withdrawing nature of  $-NO_2$  group, *o*-nitrophenol is most acidic among all the given compounds.




This is Cannizzaro reaction where one molecule gets reduced to alcohol and other gets oxidised to sodium salt of carboxylic acid. Sodium benzoate is used as a food preservative.






Mechanism :



(i) Generation of electrophile



(ii) Electrophilic substitution



Ketones do not reduce Tollens' solution therefore, X is 2-propanol.



$$4.4 \text{ g of } \text{CO}_2 \text{ will contain} = \frac{6.023 \times 10^{23}}{44} \times 4.4$$

$$= 6.023 \times 10^{22} \text{ molecules of } \text{CO}_2$$

1 molecule of  $\text{CO}_2$  contains two oxygen atoms.

$$6.023 \times 10^{22} \text{ molecules of } \text{CO}_2 \text{ will contain}$$

$$2 \times 6.023 \times 10^{22} = 1.2 \times 10^{23} \text{ oxygen atoms}$$

26. (d):  $\Delta_r H = \sum B.E.(\text{Reactants}) - \sum B.E.(\text{Products})$

$$= [B.E.(\text{H}_2) + B.E.(\text{Br}_2)] - [2B.E.(\text{HBr})]$$

$$= [433 + 192 - 2 \times 364] \text{ kJ} = -103 \text{ kJ}$$

27. (c):  $K_c = [x] [3x]^3 \dots(i)$

When concentration of  $\text{OH}^-$  ions is decreased by



$\frac{1}{4}$  times,  $K_c = [x'] \left[ \frac{3x}{4} \right]^3 \dots(ii)$

Equating eq. (i) and (ii)

$$x \times (3x)^3 = x' \left( \frac{3x}{4} \right)^3$$

$$64x = x'$$


28. (a)



Equation (ii) can be obtained by multiplying equation (i) by 2 and reversing it. Therefore,  $K_1$  and  $K_2$  are related as  $K_2 = \frac{1}{K_1^2}$ .



30. (a)

31. (a):  $\text{Cu}_2\text{S}$  containing little  $\text{FeS}$  is copper matte.



33. (d): Due to small size of fluorine atoms, electrons repel each other and hence, bond dissociation enthalpy of  $\text{F}_2$  is low.

34. (d): Because of its small size, nitrogen has a strong tendency to form  $p\pi-p\pi$  bonds. It has high electronegativity and does not have any  $d$ -orbital. Also, it has high ionisation enthalpy due to its stable half-filled electronic configuration.



37. (b): CO is a  $\pi$ -donor ligand which acts as a stronger ligand.

38. (a):  $\text{Mn}^{2+} (3d^5 4s^0) \longrightarrow 5$  unpaired electrons

$\text{Cu}^{2+} (3d^9 4s^0) \longrightarrow 1$  unpaired electron

$\text{Sc}^{2+} (3d^1 4s^0) \longrightarrow 1$  unpaired electron

$\text{Cu}^+ (3d^{10} 4s^0) \longrightarrow 0$  unpaired electron

Hence,  $\text{Mn}^{2+}$  has maximum number of unpaired electrons thus, it shows maximum paramagnetic behaviour.

39. (a):  $\text{MnO}_2 + 4\text{HCl} \longrightarrow \text{Cl}_2 + \text{MnCl}_2 + 2\text{H}_2\text{O}$

(A)

(B)

$3\text{Cl}_2 + \text{NH}_3 \longrightarrow \text{NCl}_3 + 3\text{HCl}$


(excess)

(C)

40. (a):  $\text{Mn}^{2+}$  has stable half-filled electronic configuration thus, it has high third ionisation energy while  $\text{Fe}^{2+}$  on losing one more electron will acquire this stable electronic configuration. Thus,  $\text{Fe}^{2+}$  is more prone to get oxidised to +3 oxidation state.

41. (a) 42. (a)

43. (b):  $\text{CH}_3\text{CN} \xrightarrow{\text{Reduction}} \text{CH}_3\text{CH}_2\text{NH}_2 \xrightarrow{\text{HNO}_2}$   
 Ethanenitrile (A) (B)  $\text{H}_2\text{O} + \text{N}_2 + \text{CH}_3\text{CH}_2\text{OH}$

44. (a): 

45. (b): Fehling's test is for aldehydes. In sucrose, there is no free aldehyde group but glucose in its open chain structure has free aldehyde group.

46. (d) 47. (a)

48. (d): For a given value of 'l' the permissible values of 'm' are  $-l, -(l-1), \dots, 0, \dots, (l-1), l$


Thus, for  $l = 2$ ,  $m$  cannot have a value of  $-3$ .

49. (d): In  $\text{O}_2^{2-}$  and  $\text{O}_2^-$ , the added extra electron enters the antibonding orbital which decreases the bond order. In  $\text{O}_2^+$ , the electron is removed from antibonding orbital which increases the bond order.

50. (d)

51. (b): Natural rubber has 'cis' configuration at every double bond.

52. (a): Hexamethylenediamine and adipic acid polymerise to form polyamide linkages in nylon-6, 6.



53. (a): In DNA, adenine (A) forms two hydrogen bonds with thymine (T) and guanine (G) forms three hydrogen bonds with cytosine (C).

54. (a): Equanil is a tranquilizer used in depression and hypertension.

55. (b): As the size of halide ion increases the  $\text{R}-\text{X}$  bond strength decreases and the halide ion will be a good leaving group. Thus, the reactivity order is  $\text{R}-\text{I} > \text{R}-\text{Br} > \text{R}-\text{Cl} > \text{R}-\text{F}$ .

56. (d): The  $p_x$  and  $p_y$  orbitals do not overlap thus, no bond is formed.

57. (b)


58. (b): Two miscible liquids with different boiling points can be separated by distillation through selective evaporation and condensation.

59. (c):

Element	%	No. of moles	Molar ratio	Simplest ratio
C	40	$\frac{40}{12} = 3.33$	$\frac{3.33}{3.33} = 1$	1
H	13.33	$\frac{13.33}{1} = 13.33$	$\frac{13.33}{3.33} = 4$	4
N	46.67	$\frac{46.67}{14} = 3.33$	$\frac{3.33}{3.33} = 1$	1

Thus, the empirical formula is  $\text{CH}_4\text{N}$ .

60. (b):  $\text{HNO}_3 + 2\text{H}_2\text{SO}_4 \longrightarrow \text{NO}_2^+ + 2\text{HSO}_4^- + \text{H}_3\text{O}^+$






# ADVANCED CHEMISTRY BLOC


## (GEOMETRICAL ISOMERS)

### FOR ALKENES, OXIMES AND AZO COMPOUNDS


The primary requirement for an open chain compound to show geometrical isomerism is that, the double bonded carbon must have two different substituents. This is not required for double bonded nitrogen as nitrogen is trivalent. A few examples are :




(No geometrical isomers)



(Two geometrical isomers)




(Two geometrical isomers)




(Two geometrical isomers)

### Physical properties


- If the geometrical isomers do not differ much in polarity, the one that has higher enthalpy has the higher boiling point, higher density and higher refractive index. This is the *Auwers-Skita rule*. For example,




(Higher enthalpy, higher b.pt., higher density and higher refractive index)



- If the isomers differ much in polarity, the more polar one has the higher boiling point, higher density and higher refractive index. For example,




(More polar, higher boiling point, higher density and higher refractive index)




### Stability

- In general *trans*-isomer is more stable than *cis*-isomer. The stability of isomers of butene determined from their heats of combustion is as follows :


Mukul C. Ray, Odisha



General, stability order is :



- For dihalo-ethylenes, the *cis*-form is more stable than *trans* due to forces of attraction coming from the partial charges.







- Exceptions are only di-iodo and bromo-iodo compounds, where the *trans*-form is more stable than *cis* due to steric factors.

### FOR CONJUGATED DIENES

Number of geometrical isomers for a conjugated diene depends on the number of double bonds and whether the end groups are different or identical.

- When the end groups are different :  
e.g.,  $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}=\text{CH}-\text{C}_2\text{H}_5$  has four geometrical isomers.





- When the end groups are identical :  
e.g.,  $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}=\text{CH}-\text{CH}_3$  has three geometrical isomers.

If the pattern similar to the previous example is followed then (B) and (D) become equal.

- The number of geometrical isomers can be calculated using mathematical rules :

- When the end groups are different, and  $n$  is the number of conjugated double bonds :

$$\text{No. of geometrical isomers} = 2^n$$

- When the end groups are identical and  $n$  is the number of conjugated double bonds :



$$\text{No. of geometrical isomers} = 2^{n-1} + 2^{p-1}$$

$$p = \frac{n}{2} \text{ when } n \text{ is even and}$$



$$p = \frac{(n+1)}{2} \text{ when } n \text{ is odd.}$$

### FOR CUMULATED DIENES


Geometrical isomers are observed when the number of double bonds is odd. For example,



Rings can replace such double bonds as well.




For even number of double bonds, geometrical isomers are not possible as the groups are equidistant in space.









### FOR CYCLOALKENES

*Cis*-isomers are possible for all ring sizes. Both *cis*-form and *trans*-form exist from ring size eight and onwards.



For cyclooctene, *cis*-form is more stable than *trans*.

### FOR RING COMPOUNDS CARRYING SUBSTITUENTS



mtg

## Online Test Series

Practice Part Syllabus/ Full Syllabus  
Mock Test Papers for

# JEE Main

Log on to <http://test.pcmbtoday.com>

# Boost your fundamentals with MTG's Objective series



Deep knowledge and crystal clear understanding of fundamentals is key to success. MTG's Objective series is created keeping just this insight in mind for Class XI & XII students preparing to compete in entrance exams. Put together by MTG's renowned editorial team, these unique books ensure students get just the start they need.

## HIGHLIGHTS:

- 5,255+ pages covering the latest syllabus of AIPMT and other entrance exams
- Check-Your-Grasp questions for self-assessment
- NCERT xtract from NCERT books
- Question Banks including questions from previous years' test papers (solved) of various exams like AIIMS, AIPMT, NEET, AFMC, BHU, AMU, J&K CET, UGET-Manipal, Odisha, UP-CPMT, WB JEE, Kerala PMT, etc.
- Detailed solutions to MCQs for clear understanding
- Additional information for students for exams such as AIIMS, AIPMT, AMU, etc.



Scan now with your  
smartphone or tablet\*

# YOU ASK WE ANSWER

**Do you have a question that you just can't get answered?**

Use the vast expertise of our mtg team to get to the bottom of the question. From the serious to the silly, the controversial to the trivial, the team will tackle the questions, easy and tough.

The best questions and their solutions will be printed in this column each month.

**Q1. Why only promethium is radioactive in lanthanoids and not any other lanthanoid even it is in middle of the series?**

*(Pinaki Chattopadhyay)*

**Ans.** Abundance of the elements and the number of naturally occurring isotopes vary regularly in lanthanides. Elements with even atomic number are more abundant than their neighbours with odd atomic numbers (Harkin's rule). Stability of a nucleus is related to both, the number of neutrons and the number of protons in the nucleus.

Element promethium ( $Z = 61$ ) does not occur naturally. Its absence may be explained by Mattauch's rule which states that if two adjacent elements in the periodic table have isotopes of the same mass number, one of these isotopes must be radioactive. Two nuclides that have same mass number can both be stable only if their atomic number differ by more than one. Since elements 60 and 62 have seven isotopes each, there are not many stable mass numbers available for promethium.

According to Mattauch's rule, if promethium is to have a stable isotope, it must have a mass number outside the range of 142-150.

**Q2. Why chemotherapy for cancer treatments results in the side effects like loss of hair or anaemia?**

*(Bishal Modak, West Bengal)*

**Ans.** Chemotherapy damages all rapidly dividing cells. Cancer cells divide much more often than most normal cells, so chemotherapy damages cancer cells and can destroy them.

But some types of normal cells also divide very often. This happens in tissues where there is need for a steady supply of new cells e.g., skin, hair and nails. Chemotherapy can also damage those cells and this causes side effects. It may cause hair loss all over the body e.g., eyelashes, eyebrows, etc. However, most of the times hair loss is temporary and lost hair regrow after sometime when the treatment ends.

Chemotherapy also affects the bone marrow which makes new blood cells. Hence, during chemotherapy the level of blood cells goes down.

**Q3. Photoelectric effect is quite common in metals. Can it occur in non-metals and other substances such as water?**

*(Sandhya Sharma, Mumbai)*

**Ans.** Non-metals cannot exhibit photoelectric effect because only metals have extra electrons available in their outer shells that require lower ionisation energy. These extra electrons can be easily removed by a photon provided it has energy equal to the threshold energy i.e., minimum energy required for the removal of an electron.

Some non-metals mostly in the metalloid group can exhibit weak photoelectric effect if the energy of the incoming photon is high enough like X-rays. Covalent compounds such as water are much less likely to show any photoelectric effect because all electrons are paired and are strongly held in shared electron shells.

**Q4. Can a crystalline substance change into amorphous on its own?**

*(Rohtash, U.P.)*

**Ans.** The crystalline state is relatively more stable than the amorphous state. So the Gibbs' free energy  $\Delta G$ , of a substance in the amorphous state is always higher than the Gibbs' free energy of same substance in its crystalline state. This accounts for the transformation of an amorphous solid into its crystalline state. The reverse transformation from the crystalline state to the amorphous state does not occur because it is energetically unfavourable. Transformation of amorphous substances into crystalline solids is rarely observed. This is attributed to the high viscosity of solids which allows change to occur but only very slowly. ♦♦



Minimise your efforts to  
excel in **Boards & Competitive Exams**

✓ Error-free    ✓ Easy Language    ✓ Authentic Solutions    ✓ Fully Solved by Experts

# NCERT

## EXERCISES + EXEMPLAR SOLUTIONS

A number of questions in School Examinations/Boards, Medical, Engineering Entrance Exams and Olympiads are directly picked from NCERT books. Going by Experts' and Toppers' advice, a thorough revision of NCERT textbooks will definitely widen your chances of selection in any exam.

Available for Class 6 to 12



Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!


You may find free NCERT Solutions at many places  
but why MTG is better than the rest

- Comprehensive Guide that covers entire solutions of NCERT Textbook Questions and NCERT Exemplar Books questions
- Expert answers by experienced teachers who check CBSE Papers
- Error free, Authentic solutions with the accurate information
- Complete solution for all the questions



Available at all leading book shops throughout the country.  
For more information or for help in placing your order:  
Call 0124-6601200 or email: [info@mtg.in](mailto:info@mtg.in)

# BITSAT is a little different.



Free  
Interactive  
CD with  
MTG's BITSAT  
Explorer  
Simulate the  
online testing  
experience with this  
unique interactive CD.  
Runs on both PCs  
and Macs.

## FEATURES:

- Covers all 5 subjects - Physics, Chemistry, Mathematics, English & Logical Reasoning
- Chapter-wise MCQs for practice with 1,000+ MCQs in each section
- 5 Model Test Papers to assess readiness and chances of success

There are two kinds of students out there when it comes to engineering entrance exams. Those who believe in a carpet-bombing approach - prepare for JEE and the rest of the exams get taken care of automatically. And others who are a bit focussed in their choice. MTG's BITSAT Explorer is for the latter.

Get a real-world feel of BITSAT with MTG's BITSAT Explorer. Find out what's different about BITSAT - its pattern of examinations and key success factors. Attempt the papers in given CD to check your chances of success. Either way, at Rs 525, isn't this a steal for BITSAT aspirants like you? So what are you waiting for? Order MTG's BITSAT Explorer today.



Scan now with your  
smartphone or tablet  
Application to read  
QR codes required

Available at all leading book shops throughout the country. To buy online visit [www.mtg.in](http://www.mtg.in).


For more information or for help in placing your order, call 0124-6601200 or e-mail [info@mtg.in](mailto:info@mtg.in)



# CHEMISTRY MUSING

## SOLUTION SET 34


1. (a):




2. (b): Temperature coefficient =  $\frac{k_{373 \text{ K}}}{k_{363 \text{ K}}}$

$$= \frac{(t)_{363 \text{ K}}}{(t)_{373 \text{ K}}} = \frac{300}{180} = 1.66 \quad \left( k \propto \frac{1}{t} \right)$$

3. (a): According to Baeyer's strain theory, stability decreases as the size of the ring decreases. Further, because of the presence of a double bond in a three-membered ring, cyclopropene (ii) is the least stable. Thus, the order of stability is (iv) > (iii) > (i) > (ii).




Mechanism:



5. (b): This statement governs the third law of thermodynamics.

6. (c): Compound (c) shows optical inactivity as there is no chiral C-atom and a plane of symmetry is present in the compound.



8. (c)

9. (4):  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$ ,  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  and  $[\text{CoF}_6]^{3-}$  are outer-orbital complexes having  $sp^3d^2$  hybridisation. In these complexes, due to presence of weak field ligands *i.e.*,  $\text{H}_2\text{O}$  and  $\text{F}^-$ , pairing will not occur. Hence, shows  $sp^3d^2$  hybridisation.  $[\text{Pt}(\text{NH}_3)_6]^{4+}$ ,  $[\text{Fe}(\text{CN})_6]^{4-}$ ,  $[\text{Co}(\text{CN})_6]^{3-}$  and  $[\text{Ir}(\text{NH}_3)_6]^{3+}$  are inner-orbital complexes having  $d^2sp^3$  hybridisation.

In these complexes, due to presence of strong field ligands *i.e.*,  $\text{CN}^-$  and  $\text{NH}_3$ , pairing will take place. Also Pt and Ir has high  $\Delta_o$  value, therefore pairing will occur instead of filling electrons in other orbitals because of high energy requirement.

10. (9): For B-H compound,

$$P = 0.658 \text{ atm}, V = \frac{407}{1000} \text{ litre}, T = 373 \text{ K}, w = 0.553 \text{ g}$$

$$\therefore PV = \frac{w}{M} RT; 0.658 \times \frac{407}{1000} = \frac{0.553}{M} \times 0.821 \times 373$$

$$M = 63.23 \text{ g mol}^{-1}$$

$$\therefore 100 \text{ g of compound has } 85.7 \text{ g of B}$$

$$\therefore 63.23 \text{ g of compound has } \frac{85.7 \times 63.23}{100} = 54.19 \text{ g of B}$$

$$= \frac{54.19}{10.8} \text{ g-atoms of B}$$

$$= 5 \text{ g-atoms of B} \quad [\because 10.8 \text{ g} = 1 \text{ g-atom of B}]$$


$$\therefore \text{Formula becomes } \text{B}_5\text{H}_y$$

$$\therefore 5 \times 10.8 + y = 63.23; y = 9.23 \approx 9$$

$$\therefore \text{Formula of compound is } \text{B}_5\text{H}_9, \text{ hence } y \text{ is } 9.$$



# Master Resource Books in Chemistry



## HIGHLIGHTS:

- Concise theory for competitive exams
- Illustrations with detailed solutions
- Twists and turns to learn important formulae
- Elaborate solutions to mysterious NCERT problems
- Practice assignments with pinch of hints
- Solved MCQs single and multiple option correct type, Assertion & Reason, Fill in the blanks, True or False, Comprehension, Integer & Matching types with Miscellaneous Questions

Success in **CET 2016 Now Made Easy**



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or email: info@mtg.in

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!

# CROSSWORD



Readers can send their responses at [editor@mtg.in](mailto:editor@mtg.in) or post us with complete address by 25<sup>th</sup> of every month to win exciting prizes.  
Winners' name with their valuable feedback will be published in next issue.


## ACROSS

2. Ternary acids of hydrogen, oxygen and a non-metal. (8)
4. One of the polymorphs of zinc sulphide. (8)
7. Another name of sodium aluminosilicate. (9)
9. Colour imparted to glass when it is mixed with manganese dioxide. (6)
10. Electronegativity scale based upon bond energy data. (7)
14. Nature of reversible adiabatic processes. (10)
15. A crystalline halogen. (6)
16. Nickel based superalloys containing chromium and other additives like titanium and aluminium. (7)
17. An alcohol water mixture containing 57.1% of ethylalcohol by volume. (11)
19. The acidic hydride of nitrogen. (13)
20. The total life span of a radioactive element. (8)
27. Type of isomerism exhibited by urea. (11)
28. Geometry of  $N(SiH_3)_3$ . (6)
30. A reducing agent used in vat dyeing obtained from formaldehyde sodium bisulphite derivative. (9)
12. Enzyme used to dissolve blood clots. (13)
13. Method of refining used when metal contains impurities of its own oxide. (6)
18. Another name of iron carbide. (9)
21. Complexes of NO with transition metal ions. (9)
22. Ground state of nitrenes. (7)
23. Non-protein component of enzyme. (8)
24. The difference between oxidation number of Mn in permanganate and Fe in haematite. (4)
25. Unsaturated hydrocarbons containing double bonds on adjacent or successive carbon atoms. (9)
26. A piezoelectric material. (6)
29. Can be used to put off metal fires. (4)

## DOWN

1. Linear polymer of  $\beta$ -D-glucose is which  $C_1$  of one glucose unit is connected to  $C_4$  of the other through  $\beta$ -D-glucosidic linkage. (9)
3. Solutions of two electrolytes having same concentration of common ions. (9)
5. In E-Z notation for geometrical isomers, the word that signifies symbol Z. (8)
6. Synthesis of petrol by hydrogenation of coal was invented by \_\_\_\_\_. (7)
8. A highly toxic element of group 13. (8)
11. Drug used for treatment of cancer. (5)

Cut Here





# Now, save up to Rs 2,020\*



## Subscribe to MTG magazines today.

Our 2016 offers are here. Pick the combo best suited for your needs. Fill-in the Subscription Form at the bottom and mail it to us today. If in a rush, log on to [www.mtg.in](http://www.mtg.in) now to subscribe online.

\*On cover price of ₹ 30/- each.

## About MTG's Magazines

Perfect for students who like to prepare at a steady pace, MTG's magazines – Physics For You, Chemistry Today, Mathematics Today & Biology Today – ensure you practice bit by bit, month by month, to build all-round command over key subjects. Did you know these magazines are the only source for solved test papers of all national and state level engineering and medical college entrance exams?

For JEE (Main & Advanced), NEET, PMTs, All State Level Engg. & Medical Exams

Trust of over 1 Crore readers. Since 1982.

- Practice steadily, paced month by month, with very-similar & model test papers
- Self-assessment tests for you to evaluate your readiness and confidence for the big exams
- Content put together by a team

comprising experts and members from MTG's well-experienced Editorial Board

- Stay up-to-date with important information such as examination dates, trends & changes in syllabi
- All-round skill enhancement –

confidence-building exercises, new studying techniques, time management, even advice from past IIT/PMT toppers

- **Bonus:** exposure to competition at a global level, with questions from Intl. Olympiads & Contests

## SUBSCRIPTION FORM

Please accept my subscription to:  
(Confirm your choice by ticking the appropriate boxes)

### PCMB combo

- ☐ 1 yr: ₹ 1000 (save ₹ 440) ☐ 2 yrs: ₹ 1,800 (save ₹ 1,080) ☐ 3 yrs: ₹ 2,300 (save ₹ 2,020)

### PCM combo

- ☐ 1 yr: ₹ 900 (save ₹ 180) ☐ 2 yrs: ₹ 1,500 (save ₹ 660) ☐ 3 yrs: ₹ 1,900 (save ₹ 1,340)

### PCB combo

- ☐ 1 yr: ₹ 900 (save ₹ 180) ☐ 2 yrs: ₹ 1,500 (save ₹ 660) ☐ 3 yrs: ₹ 1,900 (save ₹ 1,340)

### Individual magazines

☐ Physics ☐ Chemistry ☐ Mathematics ☐ Biology

- ☐ 1 yr: ₹ 330 (save ₹ 30) ☐ 2 yrs: ₹ 600 (save ₹ 120) ☐ 3 yrs: ₹ 775 (save ₹ 305)

Name: \_\_\_\_\_

Complete Postal Address: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Pin Code       Mobile #

Other Phone #

Email \_\_\_\_\_

Enclose Demand Draft favouring MTG Learning Media (P) Ltd, payable at New Delhi. You can also pay via Money Orders. Mail this Subscription Form to Subscription Dept., MTG Learning Media (P) Ltd, Plot 99, Sector 44, Gurgaon - 122 003 (HR).

Note: Magazines are despatched by Book-Post on 4th of every month (each magazine separately). Should you want us to send you your copies by Courier or Regd. Post instead, additional charges apply (₹ 240, ₹ 450, ₹ 600 for 1-yr, 2-yr, 3-yr subscriptions respectively).

E-mail [info@mtg.in](mailto:info@mtg.in). Visit [www.mtg.in](http://www.mtg.in) to subscribe online. Call (0)8800255334/5 for more info.